

AD _____

Award Number: DAMD17-98-1-8263

TITLE: Temporal Control of Transforming Growth Factor (TGF) -
betal Expression on Mammary Cell Multistep Transformation

PRINCIPAL INVESTIGATOR: Nancy Dumont

CONTRACTING ORGANIZATION: Vanderbilt University Medical Center
Nashville, Tennessee 37232-2103

REPORT DATE: October 2001

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20020717 028

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)**2. REPORT DATE**

October 2001

3. REPORT TYPE AND DATES COVERED

Annual Summary (1 Oct 00 - 30 Sep 01)

4. TITLE AND SUBTITLE

Temporal Control of Transforming Growth Factor (TGF) -
betal Expression on Mammary Cell Multistep Transformation

5. FUNDING NUMBERS

DAMD17-98-1-8263

6. AUTHOR(S)

Nancy Dumont

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Vanderbilt University Medical Center
Nashville, Tennessee 37232-2103

E-Mail: nancy.dumont@mcmail.vanderbilt.edu

**8. PERFORMING ORGANIZATION
REPORT NUMBER****9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)**

U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

**10. SPONSORING / MONITORING
AGENCY REPORT NUMBER****11. SUPPLEMENTARY NOTES****12a. DISTRIBUTION / AVAILABILITY STATEMENT**

Approved for Public Release; Distribution Unlimited

12b. DISTRIBUTION CODE**13. ABSTRACT (Maximum 200 Words)**

Transforming growth factor- β (TGF β) is a pleiotropic growth factor, which plays a critical role in modulating cell growth, differentiation and plasticity. There is increasing evidence that after cells lose their sensitivity to TGF β -mediated growth inhibition, autocrine TGF β signaling may potentially promote tumor progression. In order to further understand the role of TGF β in mammary transformation, the MDA-MB-231 human metastatic breast cancer cell line was used to generate stable cell lines expressing a dominant negative TGF β type II receptor (dnT β RII). Expression of dnT β RII was confirmed by affinity labeling cell surface receptors with 125 I-TGF β 1 and immunoprecipitating the affinity-labeled exogenous receptor via its HA tag. The dnT β RII was determined to be functional in that it could associate with the type I TGF β receptor and prevent it from initiating signal transduction. Results to date indicate that the basal migratory potential of dnT β RII expressing cells is impaired, and this impairment appears to be TGF β specific. Biochemical experiments are currently underway to identify the signal transducers perturbed by dnT β RII expression which impair motility, and which may, in turn, be relevant to TGF β -mediated invasion and metastases.

14. SUBJECT TERMS

Breast Cancer, Transforming growth factor beta, migration, signaling

15. NUMBER OF PAGES

41

16. PRICE CODE**17. SECURITY CLASSIFICATION
OF REPORT**

Unclassified

**18. SECURITY CLASSIFICATION
OF THIS PAGE**

Unclassified

**19. SECURITY CLASSIFICATION
OF ABSTRACT**

Unclassified

20. LIMITATION OF ABSTRACT

Unlimited

Table of Contents

Cover.....	1
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	10
Reportable Outcomes.....	11
Conclusions.....	11
References.....	12
Appendices.....	13

Introduction

The overall goal of this research was to elucidate the complex role of transforming growth factor β s (TGF β s) in sequential stages of mammary epithelial transformation, and to determine whether blocking autocrine TGF β signaling in MDA-231 human breast cancer cells would affect their metastatic phenotype *in vitro* and *in vivo*. This knowledge is of potential importance for the development of rational therapeutic approaches in human breast carcinoma.

Body

Specific Aim 1

The purpose of this Aim was to generate MMTV/tTA + tet-op/TGF β 1^{S223/225} mice. As stated in our 1999 Annual report, although MMTV/tTA mRNA was expressed in the mammary glands of the F1 bigenic animals we generated, expression levels were very low, and tet-op/TGF β 1 mRNA was not detectable. Since this may have been due to the low level of expression of the MMTV/tTA, we obtained other MMTV/rtTA founder lines (1 heterozygous male and 1 heterozygous female) with more robust MMTV expression in the mammary gland from Dr. Lewis Chodosh (University of Pennsylvania, PA). Unfortunately, the male MMTV/rtTA mouse died while being quarantined in our animal facility and the female MMTV/rtTA mouse died shortly after being released from quarantine. Consequently, breedings between these mice and our tet-op/TGF β 1 homozygous founder lines could not be set-up in order to generate the desired bigenic mice. We have since obtained two more male MMTV-rtTA mice from Dr. Chodosh. These mice are currently being quarantined in our mouse facility. Once the two male MMTV-rtTA mice have been released from quarantine (scheduled for November 5, 2001), breeding pairs will be established between the MMTV-rtTA males and the tet-op/TGF β 1 females in order to generate the bigenic mice required for our studies.

Specific Aim 2

This Aim initially proposed to study the effect of mammary TGF β 1 overexpression on different stages of breast transformation in MMTV/neu + TGF α bigenic mice. Due to the complexity of generating a mouse between these two bigenics bearing four different transgenes, two alternative approaches were proposed in our 1999 Annual Report. However, until an appropriate MMTV/tTA + tet-op/TGF β 1 bigenic mouse is generated, these studies cannot be pursued.

Specific Aim 3

This Aim initially proposed to test the effect of antisense TGF β 1 and antisense TGF β 2 on MDA-231 human breast cancer cells. However, as stated in our 2000 Annual Report, since we were unable to generate stable transfectants with sustained expression of antisense, we chose to disrupt autocrine TGF β signaling in these cells by stably expressing a dominant negative type II TGF β receptor.

Generation of MDA-MB-231 Cells Stably Expressing a dnT β RII

To generate MDA-MB-231 cells stably expressing a dnT β RII, we obtained a construct encoding a kinase inactive T β RII in which the lysine at position 277 has been mutated to arginine (dnT β RII-K277R) (Oft et al., 1998) from Martin Oft (UCSF, San Francisco, CA). Lysine 277 corresponds




Fig. 1 Generation of MDA-MB-231 Cells Stably Expressing a dnT β RII.

to an invariant lysine found in the ATP-binding site of subdomain II in all protein kinases, and even its substitution with arginine results in loss of kinase activity (Wrana et al., 1992). The control pGABE and dnT β RII encoding vectors are modified versions of the commonly used retroviral vector pBABE in which the puromycin cassette has been replaced by enhanced green fluorescent protein (GFP). In this construct, dnT β RII-K277R is HA tagged, and its expression is driven by the viral long terminal repeat (LTR), while expression of GFP is driven by the SV40 promoter (Fig. 1). Following transfection, cells expressing GFP were sorted by flow cytometry. After multiple sortings, a pool of transfectants, the majority of which expressed GFP, was obtained (Fig. 1). Clones were then isolated by sorting individual cells from the positive pool.

Verification of dnT β RII-K277R Expression in MDA-MB-231 Cells

To verify that the dnT β RII-K277R mutant was indeed expressed in the pool and clones of MDA-MB-231 cells expressing EGFP, cell surface receptors were affinity labeled with 125 I-TGF β 1 (Fig. 2). As expected, cell surface labeling of parental cells resulted in the labeling of three proteins corresponding to the endogenous type I, II, and III TGF β receptors expressed in these cells. There was little or no change in the amount of labeling observed in the control GABE clones. However, clones expressing dnT β RII-K277R, displayed a significant increase in the amount of labeled type II receptor, suggesting that the exogenous receptor was indeed expressed. A similar pattern of labeling was observed in the GABE pool from which the GABE clones were sorted, and the dnT β RII-K277R pool from which the dnT β RII-K277R clones were sorted.

In order to confirm that this increase in the labeling of the type II receptor was indeed due to expression of dnT β RII-K277R, extracts from affinity labeled cells were immunoprecipitated with an anti-HA antibody to pull down the HA-tagged dnT β RII-K277R. As shown in Fig. 3, the anti-HA antibody immunoprecipitated a labeled type II receptor in the pool and clones expressing dnT β RII-K277R, but not




Fig. 2. Expression of dnT β RII Increases Cell Surface Receptor Labeling. MDA-MB-231 parental cells as well as clones and pools stably expressing dnT β RII or the control GABE vector were affinity labeled with 2.5 ng/ml 125 I-TGF β 1 and cross-linked with BS 3 . Labeled ligand-receptor complexes were resolved by SDS-PAGE using a 3-12% gradient gel and visualized by autoradiography. Molecular weight markers are indicated on the left, and TGF β receptors are indicated on the right.




Fig. 3. Immunoprecipitation of Affinity Labeled HA-Tagged dnT β RII. Cells were affinity labeled with 2.5 ng/ml 125 I-TGF β 1, lysed and incubated with a mouse monoclonal anti-HA antibody for immunoprecipitation of HA-tagged dnT β RII. Immunoprecipitates were resolved by SDS-PAGE using a 7.5% polyacrylamide gel and visualized by autoradiography. Affinity labeled, but non-immunoprecipitated G15-5 cells were loaded as a reference (NIP).

in the control GABE pool or clones, confirming transgene expression. The type I and type III TGF β receptors appeared to co-immunoprecipitate with dnT β RII-K277R in these experiments. This was confirmed in subsequent co-immunoprecipitation experiments (see Fig. 4 below).

Evaluation of dnT β RII-K277R Function in MDA-MB-231 Cells

Having ascertained that dnT β RII-K277R was indeed expressed in the MDA-MB-231 pool and clones, we then wished to determine whether it was functional. If so, it should associate with the type I receptor, prevent the endogenous type II receptor from associating with, phosphorylating, and activating the type I receptor, and therefore prevent the type I receptor from initiating signal transduction. The ability of dnT β RII-K277R to act as a dominant negative was evaluated utilizing four different assays. First, co-immunoprecipitation experiments were performed in order to determine whether dnT β RII-K277R could in fact associate with the type I TGF β receptor (T β RI). Next, the effect of dnT β RII-K277R expression on various steps of TGF β -mediated signal transduction was examined by i) immunoblot analysis of phospho-Smad2, ii) immunofluorescence staining to monitor Smad2 translocation to the nucleus, and iii) transcriptional activation of TGF β -responsive reporter constructs.

i) Co-immunoprecipitation of T β RI with dnT β RII-K277R

The co-immunoprecipitation experiments revealed that when affinity labeled cells expressing dnT β RII-K277R were immunoprecipitated with an anti-HA antibody, a labeled protein the size of a type I receptor co-immunoprecipitated with the dnT β RII-K277R (Fig. 3 and Fig. 4, right panel – last lane). We confirmed that this was T β RI by immunoprecipitating similarly labeled cells with various TGF β superfamily type I receptor antibodies, and observed that the only antibody which immunoprecipitated a type I receptor was the T β RI antibody, ALK5 (Fig. 4). These data suggest that dnT β RII-K277R can in fact associate with the type I TGF β receptor.




Fig. 4. IP of ^{125}I -TGF β 1-Labeled Receptors. MDA-MB-231 pools expressing dnT β RII (right panel), or control GABE vector (left panel) were affinity labeled with 2.5 ng/ml ^{125}I -TGF β 1, cross-linked with BS 3 , lysed, and incubated with NRS (normal rabbit serum), polyclonal rabbit anti-sera directed against various type I receptors (ALK1,2,5,&8), the type II TGF β receptor (T β RII), or HA, as indicated. Immunoprecipitates were resolved by SDS-PAGE using a 3-12% gradient gel and visualized by autoradiography. Affinity labeled, but non-immunoprecipitated cells were loaded as a reference (NIP).

ii) Impaired TGF β -mediated Smad2 Phosphorylation in dnT β RII-K277R Clones

We next examined the effect of dnT β RII-K277R expression on TGF β -induced Smad2 phosphorylation. Immunoblot analysis of TGF β treated cell lysates using a phospho-specific Smad2 antibody revealed that while TGF β could induce phosphorylation of Smad2 in both GABE clones (G15-5 and G15-6), its ability to do so in the dnT β RII-K277R clones was impaired (Fig. 5). This impairment was not due to a decrease in total Smad2 protein, as reprobing with an antibody directed against the unphosphorylated form of Smad2 did not reveal any significant change in total Smad2 protein.




Fig. 5. **Impaired Smad2 Phosphorylation in dnT β RII Clones.** Cells were treated with 2 ng/ml TGF β 1 for the indicated times, washed, and lysed. Protein extracts (50 μ g/lane) were separated by 7.5% SDS-PAGE followed by immunoblot analysis for phospho-Smad2 and total Smad2.

iii) Impaired TGF β -mediated Nuclear Translocation of Smad2 in dnT β RII-K277R Clones

Following phosphorylation, receptor-activated Smads associate with the common mediator Smad, Smad4, and translocate to the nucleus. Therefore, next we examined the effect of dnT β RII-K277R expression on TGF β -induced Smad translocation to the nucleus by immunofluorescence (Fig. 6). In the GABE clone, in the absence of TGF β , Smad2 staining was relatively diffuse, but upon TGF β treatment for 60 min, Smad2 staining became concentrated in the nucleus. Nuclear localization of Smad2 was confirmed by staining the same cells with a nuclear dye. In contrast, in the dnT β RII-K277R clones, there was little or no change in Smad2 staining following TGF β treatment, suggesting that TGF β -mediated translocation of Smad2 to the nucleus is impaired by dnT β RII-K277R.




Fig. 6. **Impaired Smad2 Nuclear Translocation in dnT β RII Clones.** Cells were grown on glass cover-slips for 24h, serum starved for 16h, followed by treatment with 1 ng/ml TGF β 1 for 60 min. Cells were then fixed, permeabilized, and blocked with 3% milk in PBS for indirect immunofluorescence staining of Smad2 followed by nuclear staining with Hoechst. Cover-slips were mounted onto 25x75-mm microslides and fluorescent images were captured using a Princeton Instruments cooled CCD digital camera from a Zeiss Axiophot upright microscope.

iv) Impaired TGF β -mediated Transcription in dnT β RII-K277R Clones

Following translocation to the nucleus, Smads regulate gene transcription, so we examined the effect of dnT β RII-K277R expression on the ability of TGF β to induce transcription of two TGF β responsive reporter constructs, p3TP-luciferase, and pCAGA-luciferase. Each reporter construct was transiently transfected into the GABE and dnT β RII-K277R clones along with pCMV-Renilla, and luciferase activity before and after TGF β treatment was measured utilizing a dual luciferase reporter assay system. Normalized luciferase activity indicated that TGF β could induce transcription of both reporter constructs in the GABE clones (G15-5 and G15-6), but its ability to do so in the dnT β RII-K277R clones was impaired (Fig. 7).




Fig. 7. Impaired Transcription in dnT β RII Clones. Cells were transiently transfected with either (A) p3TP-Lux or (B) p(CAGA)₁₂-Lux (3 μ g/35mm dish) along with pCMV-Renilla (0.015 μ g/35mm dish). The following day, cells were split to 6 wells of a 24-well plate, treated with 1 ng/ml TGF β 1 for 16h, washed, and lysed. Firefly and Renilla luciferase activities were measured using Promega's Dual Luciferase Reporter Assay System. Luciferase activity (Y axis) is presented as the ratio of firefly to Renilla luciferase activities. Each data point represents the mean \pm SD of three wells. Fold induction of luciferase activity for each clone is indicated above the bar representing luciferase activity following TGF β 1 treatment.

Taken together, these data indicate that dnT β RII-K277R is expressed and functional in MDA-MB-231 cells in that it can associate with T β RI (ALK5), and block every step of TGF β -mediated signal transduction examined, including Smad2 phosphorylation, translocation of Smad2 to the nucleus and transcription. This system was therefore used as a model to study the molecular mechanisms by which TGF β may be contributing to tumor maintenance and progression.

Although invasion and metastasis are complicated processes that involve multiple steps, active migration of cells into the surrounding tissue is a defining characteristic of the invasive tumor (Wells, 2000). Thus, as an *in vitro* biological assay that may be relevant to the metastatic phenotype *in vivo*, we chose to examine the effect of dnT β RII-K277R expression on the motility of MDA-MB-231 cells in a wound closure assay. In this assay, a confluent monolayer of cells is wounded with a pipet tip, and the ability of the cells to migrate and close the wound is monitored by microscopy. In the GABE clones, complete wound closure was observed by 24h, while in the dnT β RII-K277R clones, although there was some closure, the wound remained open at 24h (Fig. 8). This difference in motility did not appear to be due to an effect on proliferation because when the experiment was performed in the presence of mitomycin C, a




Fig. 8. Impaired Motility in dnTβRII Clones. Confluent cell monolayers were wounded with a pipet tip. Following wounding, cell culture medium was replaced with fresh medium containing serum and wound closure was monitored by microscopy at the times indicated.




Fig 9. ALK5 Restores Motility in dnTβRII 15-2. Cells were infected with adenoviruses encoding constitutively active mutants of the BMP (ALK2^{QD}, ALK3^{QD}) or TGFβ (ALK5^{TD}) type I receptors. Uninfected cells, and cells infected with a β-GAL adenovirus were used as controls. Following infection, cell monolayers were wounded with a pipet tip, and wound closure was monitored by microscopy at the times indicated.

compound that inhibits cell division, the same results were obtained. Thus, expression of dnTβRII-K277R in MDA-MB-231 cells appears to impair their motility, independent of changes in proliferation.

Since we have data suggesting that in addition to blocking TGFβ signaling, expression of dnTβRII-K277R can also block BMP signaling in MDA-231 cells, we obtained adenoviruses encoding constitutively active mutants of TGFβ and BMP type I receptors in order to assess the relative contribution of each pathway in motility. Our results indicate that the impairment in motility observed in our cells appears to be TGFβ specific in that it could be restored by expression of a constitutively active type I TGFβ receptor (ALK5^{TD}), but not by constitutively active BMP type I receptors (ALK2^{QD} and ALK3^{QD}), Fig. 9. Consistent with this, expression of Smad6, an inhibitory Smad which preferentially blocks BMP signaling, had no effect on the migration of parental or vector control cells in wound closure assays (Fig. 10).

Effect of dnTβRII-K277R Expression on MDA-231 Tumorigenesis and Metastasis

Tumor studies in nude mice were conducted in order to evaluate the effect of dnTβRII-K277R expression on tumor formation and metastases *in vivo*. However, these experiments were not informative as the control cells expressing GFP alone, failed to form tumors. We suspect that there may have been some toxicity associated with the expression of GFP as parental cells not expressing GFP did form tumors




Fig. 10. Smad6 Expression does not affect Motility.

Cells were infected with Smad6 or control βGAL encoding adenoviruses. Following infection, Smad6 expression was verified by immunoblot analysis utilizing an anti-FLAG antibody (A), and Smad6 function was evaluated by examining its ability to block BMP2-mediated Smad1 phosphorylation (B). Despite being expressed and functional, Smad6 did not impair motility in wound closure assays (C).

following subcutaneous or orthotopic injections in nude mice. We have therefore generated additional stable cell lines expressing dnTβRII-K277R without GFP, and are currently characterizing these for use in future *in vivo* studies.

Key Research Accomplishments

Generation and characterization of MDA-231 pools and clones stably expressing dnTβRII-K277R or control vector with or without GFP.

Reportable Outcomes

Manuscripts:

Dumont N. Genetic and epigenetic contributions to colorectal cancer. *APMIS* 107: 711-722, 1999.

Dumont N and Arteaga CL. Tumor promoting effects of transforming growth factor- β . *Breast Cancer Res* 2: 125-132, 2000.

McEarchern JA, Kobie JJ, Mack V, Wu RS, Meade-Tollin L, Arteaga CL, Dumont N, Besselsen D, Seftor E, Hendrix MJC, and Akporiaye ET. Invasion and metastasis of a mammary tumor involves TGF- β signaling. *Int. J. Cancer* 91: 76-82, 2001.

Abstracts:

Dumont N and Arteaga CL. Autocrine Transforming Growth Factor- β Signaling in Mammary Tumor Cell Invasiveness. Presented at the AACR Pathobiology of Cancer Workshop in Colorado, July 2000.

Dumont N, Bakin AV, and Arteaga CL. Expression of a Kinase Inactive TGF β Type II Receptor Impairs Motility in Breast Cancer Cells. Presented at the Mammary Gland Gordon Conference in Rhode Island, June 2001.

Conclusions

1. The dnT β R_{II}-K277R protein is expressed on the surface of MDA-231 cells, as evidenced by immunoprecipitation of ¹²⁵I-TGF β 1-labeled HA-tagged T β R_{II}-K277R.
2. The dnT β R_{II}-K277R is functional in that it can associate with the type I TGF β receptor and impair various aspects of TGF β signaling including:
 - i) TGF β -mediated Smad2 phosphorylation
 - ii) TGF β -mediated translocation of Smad2 to the nucleus
 - iii) TGF β -mediated transcription of TGF β responsive reporter constructs (p3TP-luciferase and p(CAGA)₁₂-luciferase)
3. Expression of dnT β R_{II}-K277R in MDA-231 cells impairs their motility.
4. The impairment in motility observed in cells expressing dnT β R_{II}-K277R appears to be TGF- β specific in that:
 - i) Motility can be restored by expression of a constitutively active TGF β type I receptor, but not by constitutively active BMP type I receptors.
 - ii) Motility is unaffected by expression of Smad6, an inhibitory Smad that preferentially blocks BMP signaling.

Since the dnT β R_{II}-K277R is expressed and functional in MDA-MB-231 breast cancer cells, this provides an excellent model in which to study the molecular mechanisms by which TGF β may contribute to tumor maintenance and progression. We have begun to do this utilizing migration as an *in vitro* biological assay that may provide insights into the mechanisms of invasion and metastasis. Biochemical experiments are currently underway to determine whether TGF β -mediated motility is a Smad-dependent process that relies on changes in integrin expression and function. Identifying signal transducers involved in TGF β -

mediated motility of breast cancer cells may provide important information for the selection of rational targets for therapeutic interventions. This may lead to opportunities to selectively inhibit the non-desirable tumor promoting effects of TGF β while retaining its tumor suppressive effects.

References

- Oft M, Heider KH, Beug H (1998). TGFbeta signaling is necessary for carcinoma cell invasiveness and metastasis. *Curr Biol.* **8**:1243-52.
- Wells A (2000). Tumor invasion: role of growth factor-induced cell motility. *Adv Cancer Res.* **78**:31-101.
- Wrana JL, Attisano L, Carcamo J, et al. (1992). TGF beta signals through a heteromeric protein kinase receptor complex. *Cell.* **71**:1003-14.

Genetic and epigenetic contributions to colorectal cancer

Review article

NANCY DUMONT

Vanderbilt University School of Medicine, Department of Cell Biology, Nashville, Tennessee, USA

Dumont N. Genetic and epigenetic contributions to colorectal cancer. APMIS 1999;107:711–22.

Both genetic and epigenetic factors contribute to the development of colorectal cancer. Specific genetic changes in proto-oncogenes, tumor suppressor genes, and DNA mismatch repair genes have led to a genetic model of colorectal tumorigenesis. Recent data highlight the importance of the TGF- β signaling pathway in regulating the progression of colorectal cancer. The loss of the tumor suppressor activity of this pathway as well as the potentially cooperative genetic aberrations involving *APC*, *K-ras*, and *p53* are reviewed in the context of the multi-step adenoma-carcinoma sequence that characterizes the development of colorectal tumorigenesis. In addition, contributing epigenetic factors including age, diet, angiogenesis, and immune response are also discussed. Combining our knowledge of the genetic and epigenetic events implicated in this disease may allow a broader understanding of the pathogenesis of colorectal cancer and hence the design of better anti-tumor interventions.

Key words: Colorectal carcinoma; oncogene; tumor suppressor gene; mismatch repair; transforming growth factor- β ; epigenetic.

Nancy Dumont, Vanderbilt University Medical Center, 649 Medical Research Building II, Room 622, Nashville, TN 37232–6838, USA.

Like many cancers, colorectal cancer arises by acquisition of genetic alterations that result in cellular transformation. Based on these alterations, a genetic model for colorectal tumorigenesis has been proposed (1). However, in addition to genetic changes, several epigenetic factors have also been shown to contribute to the development of colorectal cancer. This paper reviews genetic as well as epigenetic contributions to colorectal cancer, both of which present opportunities as potential therapeutic targets.

GENETIC ABERRATIONS AND RELATED HISTOPATHOLOGICAL FEATURES

The development of colorectal cancer is a multi-step process involving a series of genetic changes in the colonic mucosa that lead sequentially to hyperplasia, adenoma, carcinoma, and

metastasis. Numerous genes, including proto-oncogenes, tumor suppressor genes, and DNA mismatch repair genes, have been implicated in the genesis of colon cancer. The discrete genetic changes currently perceived as fundamental to the multistep process of colorectal tumorigenesis are illustrated in Fig. 1. The tumorigenic process is initiated when a cell of the normal epithelium presumably undergoes a genetic change that conveys a selective growth advantage. This predisposes it to additional mutations, each of which confers further malignant potential, thereby leading to the clonal expansion of this cell (1). Thus, neoplasms of the colon are clonal in nature in that they arise from a single cell. Although sporadic mutations account for the majority of colorectal cancers, there are two hereditary syndromes in which a strong tendency to develop colorectal cancer is transmitted by dominant inheritance: Familial Adenomatous Polyposis (FAP) and Hereditary




Fig. 1. Genetic changes associated with colorectal tumorigenesis. This process is accelerated by MMR deficiency (see text for details). Abbreviations: LOF, loss of function; GOF, gain of function; MMR, mismatch repair. Reproduced from Kinzler & Vogelstein (2) with modifications.

Nonpolyposis Colorectal Cancer (HNPCC) (2). FAP is a syndrome in which an inherited defect in the *adenomatous polyposis coli* (*APC*) gene leads to the development of multiple benign polyps throughout the colon, some of which slowly progress to invasive lesions. Thus, defects in *APC* initiate the tumorigenic process, but additional mutations, such as those illustrated in Fig. 1, are required for tumor progression. In contrast to FAP, HNPCC is a syndrome characterized by the rapid progression of colorectal tumors due to inherited defects in DNA mismatch repair (*MMR*) genes. Although the tumors from patients with HNPCC go through a series of mutations similar to those described in Fig. 1, additional mutations, unique to HNPCC, have been described. Evidence for the role of the genes most commonly implicated in sporadic and hereditary colorectal tumors are discussed individually below.

APC

One of the earliest steps in the development of colorectal cancer is loss of function of the tumor suppressor gene, *APC*. This gene was first identified as the gene responsible for FAP

by demonstrating cosegregation of mutant *APC* alleles in affected kindreds (3, 4). However, the chromosome 5q region containing this gene is also frequently affected by loss of heterozygosity (LOH) events in colorectal adenomas and carcinomas from patients without polyposis (5–7). In fact, over 70% of sporadic colorectal cancers are believed to involve mutations in *APC* (8). Studies have shown that somatic mutations identified in sporadic tumors are similar to those observed in the germline of patients with FAP, often involving codon 1309 located in the mutation cluster region of the gene (9). These mutations have been identified in adenomas as small as 5 mm, consistent with the idea that mutations in the *APC* gene may be the initiating genetic event in many tumors (7). This is supported by the fact that excision of adenomatous polyps significantly reduces the incidence of cancer development (10). Further support for the role of *APC* in the development of polyps and colorectal cancer stems from studies of a mouse genetic model for FAP known as *Min* (for multiple intestinal neoplasia). The *Min* mutation, like those in many FAP patients, causes

premature truncation of the APC protein, and mice heterozygous for the *Min* allele develop multiple adenomatous polyps and cancers in their intestine (11).

With respect to histopathological features, *APC* mutations lead to dysplastic lesions affecting the tube-shaped epithelial foldings of the colon referred to as crypts. Foci of dysplastic aberrant crypts are the earliest identifiable lesions in colorectal tumorigenesis, and are believed to be the precursors of adenomas (12). In FAP patients, different types of *APC* mutations are associated with different clinical features despite the fact that virtually all mutations result in C-terminally truncated APC proteins (2). These differences are manifested primarily in extracolonic sites, but may also be manifested by an attenuated form of FAP in which patients develop fewer polyps (13). This phenotype is also observed in the *Min* mouse where, depending on the inbred strain carrying the *Min* allele, wide variations in polyp number are seen. Linkage analysis has demonstrated that much of the variation is due to a single locus, named *MOM-1* (for modifier of *Min*) (14), which encodes a secreted phospholipase A2 (15).

Although the exact mechanism by which *APC* mutations cause abnormal growth of colorectal epithelial cells is not clear, the fact that the majority of somatic and germline mutations in *APC* generate truncated APC proteins that lack a β -catenin-binding domain suggests that the interaction between APC and β -catenin may be important (16). Indeed, studies have shown that APC and glycogen synthase kinase 3 regulate cytoplasmic β -catenin levels by promoting its degradation (17). Inactivation of *APC* in colorectal cells allows β -catenin to accumulate and complex with T-cell factor 4, leading to activation of transcription and deregulated cell growth (18). The importance of APC and β -catenin in the development of colorectal cancer is further illustrated by the finding that β -catenin is mutated in a subset of colorectal cancers that lack somatic mutations in *APC* (19, 20). In addition to its role in regulating β -catenin, APC may also be involved in regulating apoptosis. Studies have shown that expression of wild-type APC in colorectal epithelial cells with *APC* mutations results in cell death (21). Thus, this may be another mechanism by which inactivation of *APC* leads to deregulated cell growth.

K-ras

Although the three *ras* genes, *K-ras*, *H-ras*, and *N-ras*, are highly homologous, and believed to be expressed at relatively equivalent levels in the colonic mucosa, only *K-ras* plays a significant role in the development of colorectal cancer. Mutations in *K-ras* can be identified in 50% of colorectal cancers (22), and occur most frequently in codon 12, with fewer mutations found at codons 13 and 61 (23). Because *K-ras* is an oncogene, mutation of one allele is enough to produce an effect. These mutations affect the ability of p21ras to interact with the ras GTPase-activating protein, causing p21ras to remain in the active GTP-bound state. As a result, the growth and differentiation signal transduction pathways that include p21ras are constitutively activated, leading to a continually growth-stimulated state.

The frequency of *ras* mutations appears to be correlated with two histopathological features: increased tumor size and dysplasia. Studies have shown that 50% of adenomas greater than 1 cm in diameter harbor *K-ras* mutations compared to only 10% of adenomas less than 1 cm (22). When adenomas are distinguished from one another with respect to the degree of dysplasia, *ras* mutations are more prevalent in tumors with increased dysplasia (24). The higher prevalence of *ras* mutations in later stage adenomas and carcinomas suggests that these mutations may arise in one cell of a small preexisting adenoma causing it to progress to a larger and more dysplastic adenoma, with greater risk of subsequent progression to cancer. This is consistent with the fact that hyperplastic cells containing mutant *ras* genes, unlike their dysplastic counterparts with mutant *APC* genes, have little or no potential to form clinically important tumors and may eventually regress through apoptosis (25).

DCC, Smad4, and the tumor suppressor activity of the TGF- β pathway

LOH affecting the long arm of chromosome 18 can be detected in more than 70% of primary colorectal cancers, in about 50% of advanced adenomas, and infrequently in earlier stage adenomas, suggesting that loss in this region is a relatively late event (22). Such losses are also correlated with greater mortality and increased propensity for metastatic spread (26, 27). Ef-

forts to identify a candidate tumor suppressor gene from 18q led to the discovery of a gene termed *DCC* (for *deleted in colorectal cancer*) (28). The *DCC* gene encodes a transmembrane protein of the immunoglobulin superfamily. The predicted structural similarity of *DCC* to the N-CAM family of cell-surface adhesion molecules suggested that it might function in differentiation pathways and cell fate determination through cell-cell and/or cell-extracellular matrix interactions (29). Therefore, it was hypothesized that loss of cell-cell contact might explain the enhanced metastasis observed in patients with loss of *DCC* (29). However, more recent studies have shown that inactivation of the murine *DCC* gene does not affect the proliferation or differentiation of intestinal epithelial cells, nor does it affect the morphogenesis of colonic crypts and villi (30). Moreover, introduction of the null *DCC* allele into the germ line of the *Min* mouse does not accelerate the progression of, or modify the phenotype of polyps initiated in the *Min* mice (30). Instead, the phenotype of mice lacking a functional *DCC* gene resembles that of netrin-1-deficient mice, with defects in axonal projections and brain development (30). These findings fail to support a tumor suppressor function for *DCC* in the development of colorectal cancer, and are inconsistent with studies in which reduction or loss of *DCC* RNA has been observed in cell lines or xenografts derived from human colon carcinomas (28, 31).

The discrepancy between these results may be due to differences in the pathogenesis of colorectal cancer between mice and humans. Alternatively, LOH of 18q21 may not only affect the *DCC* gene, but neighboring genes as well, one or more of which may be the target of inactivation during colon tumor progression (30). Indeed, other candidate tumor suppressor genes, including *Smad2* and *Smad4*, have been identified on chromosome 18q21 (32, 33). Both *Smad2* and *Smad4* belong to the *SMAD* gene family involved in the signal transduction pathways activated through the TGF- β family of receptors (34). The TGF- β s are important regulators of cell growth and differentiation (35). Escape from the growth regulatory effects of TGF- β s is common among many different cancers (35, 36), including colorectal cancer (37). Recent studies have shown that inactivation of the *Smad4* gene in *APC* ^{Δ 716} knockout mice,

which have a phenotype similar to that of the *Min* mice, results in malignant progression of the intestinal tumors at a much earlier stage than that observed in the simple *APC* ^{Δ 716} knockout mice (38). Mice heterozygous only for the *Smad4* knockout show no apparent tumor phenotype, indicating that the *Smad4* gene is a suppressor of tumor progression, but not of tumor initiation (38). The fact that inactivation of *Smad4* in *APC* ^{Δ 716} knockout mice enhances tumor progression, while inactivation of *DCC* in a similar mouse model does not, suggests that the tumor suppressor gene associated with LOH on 18q21 is more likely to be *Smad4*. In support of this, mutations in the *Smad4* gene have been identified in human colorectal cancers in vivo (33, 39, 40), and in familial juvenile polyposis, which, like FAP, is a syndrome characterized by a predisposition to hamartomatous polyps and gastrointestinal cancer (41).

A role for *Smad4* in the malignant progression of colorectal tumors is consistent with previous findings that have implicated other members of the TGF- β pathway in colorectal tumorigenesis (summarized in Fig. 2). For example, as previously mentioned, *Smad2* is a candidate tumor suppressor gene also located on 18q21. Inactivating missense mutations and deletions of the *Smad2* gene have been detected in sporadic colorectal carcinomas (32, 42). In addition to *Smad2* and *Smad4*, *Smad3* is another member of the *SMAD* family of proteins involved in mediating TGF- β signaling. Although mutations in the *Smad3* gene have not yet been detected in human colorectal cancer (43, 44), LOH of *Smad3* has been detected in 2 (1 sporadic and 1 HNPCC) of 17 colorectal cancers examined (44), and a recent study has reported that inactivation of the *Smad3* gene in mice leads to the development of metastatic colorectal cancer (45). Upstream of the *SMAD* signaling proteins, mutations in the TGF- β type II receptor (TGF- β RII) have been associated with microsatellite instability in both colon cancers and colorectal carcinoma cell lines (46–48). Restoration of TGF- β RII expression by gene transfection has been found to reverse the transformed phenotype (49). There is also evidence that transfection of TGF- β -responsive human colon carcinoma cells with a TGF- β 1 antisense expression vector increases their tumorigenicity (50). These data taken together highlight the im-

portance of the TGF- β signaling pathway in regulating the progression of colorectal cancer.

p53

The chromosomal region most frequently affected by LOH in colorectal cancers is 17p (51), a region which includes the tumor suppressor gene, *p53*. Sequence analysis of the remaining *p53* allele from a large number of colorectal carcinomas in which 17p was affected by LOH revealed that missense mutations were present in over 80% of the cases (52, 53), indicating that loss of *p53* function is an important step in the development of colorectal cancer. Although *p53* mutations are extremely prevalent in many advanced colorectal cancers, they are

much less frequent in earlier stages of carcinogenesis, and rare in both adenomas and polyps (22). Moreover, patients with germline mutations in *p53* do not develop polyposis, nor are they at risk for developing colorectal cancer (54). This suggests that, although *p53* plays an important role in colorectal carcinogenesis, unlike *APC* it cannot initiate the process. Likewise, *p53*-deficient mice are prone to develop cancers, but an initiating event is required for tumor development (55, 56). Moreover, the incidence of colorectal cancer in these animals is low, indicating that loss of *p53* function alone does not cause transformation (57). This is consistent with the role of *p53* as a cell-cycle checkpoint regulator (58). Abrogation of either *p53*-




Fig. 2. The TGF- β signaling pathway and its role in colorectal tumorigenesis. TGF- β (represented as a circle) elicits its effects through binding to specific cell-surface receptors denoted type I (RI) and type II (RII) TGF- β receptors, both of which are transmembrane serine/threonine kinases. TGF- β binds directly to RII, which is a constitutively active kinase. The ligand-bound RII then recognizes RI, leading to the formation of a heteromeric complex, allowing RII to phosphorylate and thereby activate RI. The activated RI phosphorylates Smad-2 or Smad-3, which then associates with Smad-4. This complex translocates to the nucleus where it can initiate gene transcription. TGF- β , RII, Smad-2, Smad-3, and Smad-4 have all been implicated as possible tumor suppressors in colorectal cancer, as summarized in the table above (see text for details). Abbreviations: MSI, microsatellite instability; LOH, loss of heterozygosity. Signaling pathway reproduced with modifications from Massague (34).

dependent cell-cycle arrest or apoptosis could allow cells that have incurred mutations and are not fit to progress through the cell cycle to do so anyway, resulting in tumor progression.

With respect to clinical features, the presence of mutant p53, as determined by increased immunohistochemical staining or by analysis of gene sequence, is correlated with poor survival, as well as increased cancer recurrence rates (59, 60).

MMR genes

DNA mismatch repair plays a prominent role in the correction of replicative mismatches which escape DNA polymerase proofreading. Three genes, *MutS*, *MutL*, and *MutH*, are central to the correction of replication errors in *E. coli* (61). In humans, germline mutations in the *MutS* homologue, *hMSH2* (62, 63), or in any of the three *MutL* homologues, *hMLH1*, *hPMS1*, and *hPMS2* (64–66), have been identified as being responsible for HNPCC. These reports suggest that mutations in *hMSH2* and *hMLH1* together account for the majority of HNPCC, while mutations in *hPMS1* and *hPMS2* are less frequently observed. A characteristic feature of tumors arising in individuals with HNPCC is the presence of microsatellite instability. Microsatellites are regions consisting of single, dinucleotide, or trinucleotide repeats that are widely distributed throughout the genome. These sequences are prone to replication errors due to their repetitive structure, which favors slippage during replication. Thus, when *MMR* genes are mutated, errors arising during DNA replication are less efficiently corrected, resulting in a replication error-prone (RER+) phenotype. Affected cells accumulate errors (mutations) at a much greater rate than normal cells (67). This is manifested clinically by a much faster progression of the disease. In contrast to sporadic colorectal cancers, which may take 10 to 15 years to develop, patients with HNPCC have been found to develop cancers within 2 years after a normal colonoscopy (68).

Microsatellite instability is not only a characteristic feature of HNPCC tumors, it is also observed in a subset (about 17%) of sporadic colorectal cancers (69–73). Sporadic tumors with microsatellite instability share common clinical and histopathological features with HNPCC tumors. They are usually located in the proximal

colon (71–74), are associated with extracellular mucin production (73), poor differentiation (71–73), and diploidy (69, 72). In addition, they are less likely to have LOH at known tumor suppressor gene loci on chromosomes 5q, 18q, and 17p (74). On the other hand, inactivation of TGF- β RII due to frameshift mutations within coding microsatellite sequences occurs frequently (47, 48). Although defects in the same *MMR* genes that are affected in HNPCC have been identified in sporadic colorectal cancers with microsatellite instability (75), substantial differences in the nature and incidence of these mutations have been reported, suggesting that the molecular mechanisms underlying instability in the sporadic cases differ from those in HNPCC (2, 76). However, mutations in *hMSH3* and *hMSH6*, two other *MutS* genes initially identified in sporadic colorectal tumors with microsatellite instability (77, 78), have recently been reported in HNPCC patients. This suggests that the molecular mechanisms underlying instability may be similar in tumors with microsatellite instability regardless of their sporadic or hereditary nature (79).

EPIGENETIC FACTORS THAT CONTRIBUTE TO COLORECTAL CANCER

Age and diet. Consistent with the multi-hit hypothesis for the development of cancer, the incidence of colorectal cancer increases with age, as greater numbers of mutations are accumulated with time (80, 81). Although genetic aberrations and hereditary disorders play a critical role in the development of colorectal cancers (section above), the fact that the incidence varies tremendously according to geography suggests that other factors must also be involved (82). One of the major differences in lifestyle between cultures (which may explain geographic differences in colorectal cancer incidence) is diet (83). A number of dietary components have been implicated as possible factors in the development of colorectal cancer. The major factors include overall caloric intake, fat content of the diet, and fiber intake. The number of calories consumed per day and the fat content in the diet have been consistently positively correlated with the

risk of developing colorectal cancer (80). The mechanism by which a high-fat diet enhances tumor formation may be related to high fecal bile acid levels which stimulate mucosal epithelial proliferation. In contrast, vegetable and fiber consumption seem to have a protective effect (84, 85). Fiber may work either by increasing the stool transit time and thereby decreasing contact with fecal contents, by binding luminal toxic compounds, or by providing fuel for colonic bacteria that produce short-chain fatty acids that may inhibit proliferation and promote apoptosis (86). Consistent with the protective effects of dietary fiber, the intake of selenium, an essential trace mineral found in cereal grains, has also been found to be inversely correlated with the incidence of colorectal cancer (87–89). Since selenium is a cofactor for glutathione peroxidase, which participates in preventing free radical damage to tissues, part of selenium's protective effect may be due to a reduction in free radical damage.

Angiogenesis. In addition to the risk factors associated with lifestyle, physiological processes such as angiogenesis also contribute to the pathogenesis of colorectal cancer. Studies have shown that in order for a tumor to grow beyond a few millimeters in diameter, the formation of new blood vessels is required to provide nutrients and a means of eliminating metabolic waste products (90). As tumors get larger, the center of the tumor often becomes hypoxic due to inadequate vascularization, leading to cell death within the hypoxic center. Because hypoxia can induce apoptosis in a p53-dependent manner (91), low oxygen conditions can provide a selective advantage for cells carrying mutations in p53, allowing escape from apoptosis. This may be particularly important in colorectal cancer because p53 mutations are prevalent, and occur late in the adenoma-carcinoma sequence. Thus, by escaping apoptosis, tumor cells bearing p53 mutations within a colorectal carcinoma retain their proliferative capacity, thereby promoting tumor expansion. In addition, since p53 expression results in the secretion of inhibitors of angiogenesis (92, 93), selection of p53 mutations by hypoxic conditions could lead to loss of expression of antiangiogenic factors, allowing growth of new blood vessels, thereby favoring further expansion of the tumor.

Angiogenesis is not only essential for the expansion of the primary tumor, it is also required for the establishment and growth of metastases at distant sites. In fact, there is evidence that systemic suppression of angiogenesis can maintain micrometastases dormant as a result of a balance between proliferation and apoptosis (94). Therefore, the degree of angiogenesis may be an important factor in determining tumor behavior and the propensity to metastasize. Indeed, in patients with colorectal cancer, angiogenesis has been correlated with a higher recurrence rate and diminished survival (95).

Immune response. Another epigenetic factor important in the development of colorectal cancer is ineffective immune response. Unlike virally or chemically induced tumors, spontaneous tumors, such as those arising in the colon, elicit a weak immune response. Although mutant proteins encoded by oncogenes (*K-ras*) or tumor suppressor genes (*APC*, *DCC*, *Smad4*, *Smad2*, *tgf- β RII*, *tgf- β I*, *p53*) that have undergone mutations can be recognized as tumor-specific antigens, their recognition delivers only one of the two signals required for T-cell-mediated immunity. This is due to the fact that very few professional antigen-presenting cells (dendritic cells, macrophages, and B-cells) are present in colorectal tumors (96). Thus, most of the mutant proteins are processed and presented as antigenic peptides bound to class I MHC on the surface of colon cancer cells. Since these cells are not professional antigen-presenting cells, they lack the costimulatory signal (B7) that must be recognized by T-cells in order to elicit an immune response. Recognition of foreign antigen in the absence of a costimulatory signal leads to T-cell anergy. Consequently, although a large number of T-lymphocytes have been identified in primary colorectal tumors (97), T-cell-mediated immunity is ineffective in eliminating tumor cells. In contrast, natural killer cells have a spontaneous cytotoxic capacity against tumor cells. However, these cells are either not found or are only present in low numbers in colorectal cancers (96, 98). In addition, studies have shown that MHC expression on professional antigen-presenting cells within colorectal tumors is often lost, thereby further compromising immunity against tumors (99).

Related to immune function, epidemio-

logical studies have shown that chronic use of aspirin results in a reduced risk of colorectal cancer (100), and that treatment of FAP patients with non-steroidal anti-inflammatory drugs (NSAIDs) results in regression of rectal polyps (101). These studies suggest that inflammation may contribute to the development of colorectal cancer. NSAIDs mediate their effects by inhibiting two enzymes, COX-1 and COX-2, which are responsible for eicosanoid synthesis. Analysis of COX-2 mRNA in colon cancers and adenomatous polyps revealed increases of 86% and 43%, respectively, compared to in normal mucosa from the same patients (102). Consistent with elevated COX-2 levels, prostaglandin E2 levels have also been found to be elevated in colon cancers and polyps (103, 104). Further support for the role of COX-2 in the development of colorectal cancer has been obtained using an *APC* knockout mouse with a phenotype similar to that of the *Min* mouse. Inactivation of the *COX-2* gene in these mice resulted in an 86% decrease in intestinal and colonic polyps compared to control animals (105). In addition, when animals heterozygous for *APC*, but having normal *COX-2*, were treated with a novel specific inhibitor of COX-2 enzyme activity, polyp numbers decreased by 50–60% compared to a 26% decrease when they were treated with the NSAID sulindac (105). Moreover, feeding a selective COX-2 inhibitor to rats has been reported to completely prevent the development of chemically induced colon tumors in 93% of animals (106). These data suggest a direct role of COX-2 in the development of colorectal cancer.

There is evidence that the mechanism by which COX-2 overexpression contributes to colorectal cancer involves a decrease in apoptosis due to induction of Bcl-2 by prostaglandin E2, rather than an increase in inflammation, as initially suggested (107, 108). In addition, COX-2-overexpressing colon cancer cells have been shown to produce high levels of angiogenic factors, which stimulate both endothelial migration and tube formation (109). Thus, COX-2 may also act as a tumor promoter via stimulation of angiogenesis. This is consistent with a previous study which has shown that the antitumor effect of the COX inhibitor diclofenac was due to an antiangiogenic effect (110).

CONCLUSION

Since many genetic aberrations contribute to the development of colorectal cancer, each affected gene could potentially be a good therapeutic target. Identification of these genetic aberrations has provided an opportunity to use different strategies to deliver normal copies of defective tumor suppressor genes to affected tissues, or to inactivate oncogenes. Although these avenues hold promise, not all of the genetic aberrations described above occur in all individuals with colorectal cancer. In contrast, epigenetic factors do indeed contribute to the development of all colorectal tumors and thereby provide additional opportunities for intervention. Therapies targeting both genetic and epigenetic aberrations could be combined to provide a broader and more effective therapy for patients.

I would like to thank Dr. Lynn Matrisian for critically reviewing this manuscript and Dr. Anne Lenferink for her assistance in preparing the figures. This work is supported in part by a U.S. Army Breast Cancer Pre-doctoral Training Award.

REFERENCES

1. Fearon ER, Vogelstein B. A genetic model for colorectal tumorigenesis. *Cell* 1990;61:759–67.
2. Kinzler KW, Vogelstein B. Lessons from hereditary colorectal cancer. *Cell* 1996;87:159–70.
3. Groden J, Thliveris A, Samowitz W, Carlson M, Gelbert L, Albertsen H, et al. Identification and characterization of the familial adenomatous polyposis coli gene. *Cell* 1991;66:589–600.
4. Nishisho I, Nakamura Y, Miyoshi Y, Miki Y, Ando H, Horii A, et al. Mutations of chromosome 5q21 genes in FAP and colorectal cancer patients. *Science* 1991;253:665–9.
5. Solomon E, Voss R, Hall V, Bodmer WF, Jass JR, Jeffreys AJ, et al. Chromosome 5 allele loss in human colorectal carcinomas. *Nature* 1987;328:616–9.
6. Ashton-Rickardt PG, Dunlop MG, Nakamura Y, Morris RG, Purdie CA, Steel CM, et al. High frequency of APC loss in sporadic colorectal carcinoma due to breaks clustered in 5q21–22. *Oncogene* 1989;4:1169–74.
7. Powell SM, Zilz N, Beazer-Barclay Y, Bryan TM, Hamilton SR, Thibodeau SN, et al. *APC* mutations occur early during colorectal tumorigenesis. *Nature* 1992;359:235–7.
8. Howe JR, Guillem JG. The genetics of colorec-

- tal cancer. *Surg Clin North Am* 1997;77:175-95.
9. Miyoshi Y, Nagase H, Ando H, Horii A, Ichii S, Nakatsuru S, et al. Somatic mutations of the APC gene in colorectal tumors: mutation cluster region in the APC gene. *Hum Mol Genet* 1992;1:229-33.
 10. Murakami R, Tsukuma H, Kanamori S, Imanishi K, Otani T, Nakanishi K, et al. Natural history of colorectal polyps and the effect of polypectomy on occurrence of subsequent cancer. *Int J Cancer* 1990;46:159-64.
 11. Su LK, Kinzler KW, Vogelstein B, Preisinger AC, Moser AR, Luongo C, et al. Multiple intestinal neoplasia caused by a mutation in the murine homolog of the APC gene. *Science* 1992;256:668-70.
 12. Pretlow TP, Barrow BJ, Ashton WS, O'Riordan MA, Pretlow TG, Juncisek JA, et al. Aberrant crypts: putative preneoplastic foci in human colonic mucosa. *Cancer Res* 1991;51:1564-7.
 13. Spirio L, Olschwang S, Groden J, Robertson M, Samowitz W, Joslyn G, et al. Alleles of the APC gene: an attenuated form of familial polyposis. *Cell* 1993;75:951-7.
 14. Dietrich WF, Lander ES, Smith JS, Moser AR, Gould KA, Luongo C, et al. Genetic identification of Mom-1, a major modifier locus affecting Min-induced intestinal neoplasia in the mouse. *Cell* 1993;75:631-9.
 15. MacPhee M, Chepenik K, Liddell R, Nelson K, Siracusa L, Buchberg A. The secretory phospholipase A2 gene is a candidate for the *MOM1* locus, a major modifier of APC^{Min}-induced intestinal neoplasia. *Cell* 1995;81:957-66.
 16. Nakamura Y. The role of the adenomatous polyposis coli (APC) gene in human cancers. *Adv Cancer Res* 1993;62:65-87.
 17. Rubinfeld B, Albert I, Porfiri E, Fiol C, Munemitsu S, Polakis P. Binding of GSK-3 β to the APC- β -catenin complex and regulation of complex assembly. *Science* 1996;272:1023-5.
 18. Korinek V, Barker N, Morin PJ, van Wichen D, de Weger R, Kinzler KW, et al. Constitutive transcriptional activation by a β -catenin-Tcf complex in APC^{-/-} colon carcinoma. *Science* 1997;275:1784-7.
 19. Morin PJ, Sparks AB, Korinek V, Barker N, Clevers H, Vogelstein B, et al. Activation of β -catenin-Tcf signaling in colon cancer by mutations in β -catenin or APC. *Science* 1997;275:1787-90.
 20. Iwao K, Nakamori S, Kameyama M, Imaoka S, Kinoshita M, Fukui T, et al. Activation of the β -catenin gene by interstitial deletions involving exon 3 in primary colorectal carcinomas without adenomatous polyposis coli mutations. *Cancer Res* 1998;58:1021-6.
 21. Morin PJ, Vogelstein B, Kinzler KW. Apoptosis and APC in colorectal tumorigenesis. *Proc Natl Acad Sci USA* 1996;93:7950-4.
 22. Vogelstein B, Fearon ER, Hamilton SR, Kern SE, Preisinger AC, Leppert M, et al. Genetic alterations during colorectal tumor development. *N Engl J Med* 1988;319:525-32.
 23. Bos JL, Fearon ER, Hamilton SR, Verlaan-de Vries M, van Boom JH, van der Eb AJ, et al. Prevalence of ras gene mutations in human colorectal cancers. *Nature* 1987;327:293-7.
 24. Miyaki M, Seki M, Okamoto M, Yamanaka A, Maeda Y, Tanaka K, et al. Genetic changes and histopathological types in colorectal tumors from patients with adenomatous polyposis. *Cancer Res* 1990;50:7166-73.
 25. Shpitz BHK, Medline A, Bruce WR, Bull SB, Gallinger S, Stern H. Natural history of aberrant crypt foci. *Dis Colon Rectum* 1996;39:763-7.
 26. Kern SE, Fearon ER, Tersmette KW, Enterline JP, Leppert M, Nakamura Y, et al. Clinical and pathological associations with allelic loss in colorectal carcinoma. *JAMA* 1989;261:3099-103.
 27. Jen J, Kim H, Piantadosi S, Liu ZF, Levitt RC, Sistonen P, et al. Allelic loss of chromosome 18q and prognosis in colorectal cancer. *N Engl J Med* 1994;331:213-21.
 28. Fearon ER, Cho KR, Nigro JM, Kern SE, Simons JW, Ruppert JM, et al. Identification of a chromosome 18q gene that is altered in colorectal cancers. *Science* 1990;247:49-56.
 29. Fearon ER, Ekstrand BC, Hu G, Pierceall WE, Reale MA, Bigner SH. Studies of the deleted in colorectal cancer gene in normal and neoplastic tissues. *Cold Spring Harbor Symp Quant Biol* 1994;59:637-43.
 30. Fazeli A, Dickinson SL, Hermiston ML, Tighe RV, Steen RG, Small CG, et al. Phenotype of mice lacking functional *Deleted in colorectal cancer (Dcc)* gene. *Nature* 1997;386:796-804.
 31. Kikuchi-Yanoshita R, Konishi M, Fukunari H, Tanaka K, Miyaki M. Loss of expression of the DCC gene during progression of colorectal carcinomas in familial adenomatous polyposis and non-familial adenomatous polyposis patients. *Cancer Res* 1992;52:3801-3.
 32. Eppert K, Scherer SW, Ozcelik H, Pirone R, Hoodless P, Kim H, et al. MADR2 maps to 18q21 and encodes a TGF β -regulated MAD-related protein that is functionally mutated in colorectal carcinoma. *Cell* 1996;86:543-52.
 33. Thiagalingam S, Lengauer C, Leach FS, Schutte M, Hahn SA, Overhauser J, et al. Evaluation of candidate tumor suppressor genes on chromosome 18 in colorectal cancers. *Nat Genet* 1996;13:343-6.
 34. Massague J. TGF- β signal transduction. *Annu Rev Biochem* 1998;67:753-91.
 35. Massague J. The transforming growth factor-

- beta family. *Annu Rev Cell Biol* 1990;6:597-641.
36. Filmus J, Kerbel RS. Development of resistance mechanisms to the growth-inhibitory effects of transforming growth factor- β during tumor progression. *Curr Opin Oncol* 1993;5:123-9.
 37. Hoosein NM, McKnight MK, Levine AE, Mulder KM, Childress KE, Brattain DE, et al. Differential sensitivity of subclasses of human colon carcinoma cell lines to the growth inhibitory effects of transforming growth factor- β 1. *Exp Cell Res* 1989;181:442-53.
 38. Takaku K, Oshima M, Miyoshi H, Matsui M, Seldin MF, Taketo MM. Intestinal tumorigenesis in compound mutant mice of both *Dpc4* (*Smad4*) and *Apc* genes. *Cell* 1998;92:645-56.
 39. Takagi Y, Kohmura H, Futamura M, Kida H, Tanemura H, Shimokawa K, et al. Somatic alterations of the DPC4 gene in human colorectal cancers in vivo. *Gastroenterology* 1996;111:1369-72.
 40. MacGrogan D, Pegram M, Slamon D, Bookstein R. Comparative mutational analysis of DPC4 (*Smad4*) in prostatic and colorectal carcinomas. *Oncogene* 1997;15:1111-4.
 41. Howe JR, Roth S, Ringold JC, Summers RW, Jarvinen HJ, Sistonen P, et al. Mutations in the SMAD4/DPC4 gene in juvenile polyposis. *Science* 1998;280:1086-8.
 42. Riggins GJ, Thiagalingam S, Rozenblum E, Weinstein CL, Kern SE, Hamilton SR, et al. *Mad*-related genes in the human. *Nat Genet* 1996;13:347-9.
 43. Riggins GJ, Kinzler KW, Vogelstein B, Thiagalingam S. Frequency of *Smad* gene mutations in human cancers. *Cancer Res* 1997;57:2578-80.
 44. Arai T, Akiyama Y, Okabe S, Ando M, Endo M, Yuasa Y. Genomic structure of the human *Smad3* gene and its infrequent alterations in colorectal cancers. *Cancer Lett* 1998;122:157-63.
 45. Zhu Y, Richardson JA, Parada LF, Graff JM. *Smad3* mutant mice develop metastatic colorectal cancer. *Cell* 1998;94:703-14.
 46. Lu SL, Akiyama Y, Nagasaki H, Saitoh K, Yuasa Y. Mutations of the transforming growth factor- β type II receptor gene and genomic instability in hereditary nonpolyposis colorectal cancer. *Biochem Biophys Res Comm* 1995;216:452-7.
 47. Markowitz S, Wang J, Myeroff L, Parsons R, Sun L, Lutterbaugh J, et al. Inactivation of the type II TGF- β receptor in colon cancer cells with microsatellite instability. *Science* 1995;268:1336-8.
 48. Parsons R, Myeroff LL, Liu B, Willson JK, Markowitz SD, Kinzler KW, et al. Microsatellite instability and mutations of the TGF- β type II receptor gene in colorectal cancer. *Cancer Res* 1995;55:5548-50.
 49. Wang J, Sun L, Myeroff L, Wang X, Gentry LE, Yang J, et al. Demonstration that mutation of the type II TGF- β receptor inactivates its tumor suppressor activity in replication error-positive colon carcinoma cells. *J Biol Chem* 1995;270:22044-9.
 50. Wu SP, Theodorescu D, Kerbel RS, Willson JK, Mulder KM, Humphrey LE, et al. TGF- β 1 is an autocrine-negative growth regulator of human colon carcinoma FET cells in vivo as revealed by transfection of an anti-sense expression vector. *J Cell Biol* 1992;116:187-96.
 51. Vogelstein B, Fearon ER, Kern SE, Hamilton SR, Preisinger AC, Nakamura Y, et al. Allelotype of colorectal carcinomas. *Science* 1989;244:207-11.
 52. Baker SJ, Preisinger AC, Jessup JM, Paraskeva C, Markowitz S, Willson JK, et al. p53 gene mutations occur in combination with 17p allelic deletions as late events in colorectal tumorigenesis. *Cancer Res* 1990;50:7717-22.
 53. Greenblatt MS, Bennett WP, Hollstein M, Harris CC. Mutations in the p53 tumor suppressor gene: clues to cancer etiology and molecular pathogenesis. *Cancer Res* 1994;54:4855-78.
 54. Garber JE, Goldstein AM, Kantor AF, Dreyfus MG, Fraumeni JF Jr, Li FP. Follow-up study of twenty-four families with Li-Fraumeni syndrome. *Cancer Res* 1991;51:6094-7.
 55. Donehower LA, Harvey M, Slagle BL, McArthur MJ, Montgomery CA Jr, Butel JS, et al. Mice deficient in p53 are developmentally normal but susceptible to spontaneous tumors. *Nature* 1992;356:215-21.
 56. Harvey M, Sands AT, Weiss RS, Hegi ME, Wiseman RW, Pantazis P, et al. In vitro growth characteristics of embryo fibroblasts isolated from p53-deficient mice. *Oncogene* 1993;8:2457-67.
 57. Jacks T, Remington L, Williams BO, Schmitt EM, Halachmi S, Bronson RT, et al. Tumor spectrum analysis in p53-mutant-mice. *Curr Biol* 1994;4:1-7.
 58. Sherr CJ. Cancer cell cycles. *Science* 1996;274:1672-7.
 59. Auvinen A, Isola J, Visakorpi T, Koivula T, Virtanen S, Hakama M. Overexpression of p53 and long-term survival in colon carcinoma. *Br J Cancer* 1994;70:293-6.
 60. Hamelin R, Laurent-Puig P, Olschwang S, Jeco N, Asselain B, Remvikos Y, et al. Association of p53 mutations with short survival in colorectal cancer. *Gastroenterology* 1994;106:42-8.
 61. Modrich P. Mechanisms and biological effects of mismatch repair. *Annu Rev Genet* 1991;25:229-53.
 62. Fishel R, Lescoe MK, Rao MR, Copeland NG,

- Jenkins NA, Garber J, et al. The human mutator gene homolog *MSH2* and its association with hereditary nonpolyposis colon cancer. *Cell* 1993;75:1027-38.
63. Leach FS, Nicolaides NC, Papadopoulos N, Liu B, Jen J, Parsons R, Peltomäki P, et al. Mutations of a *mutS* homolog in hereditary nonpolyposis colorectal cancer. *Cell* 1993;75:1215-25.
 64. Bronner CE, Baker SM, Morrison PT, Warren G, Smith LG, Lescoe MK, et al. Mutation in the DNA mismatch repair gene homologue *hMLH1* is associated with hereditary nonpolyposis colon cancer. *Nature* 1994;368:258-61.
 65. Papadopoulos N, Nicolaides NC, Wei YF, Ruben SM, Carter KC, Rosen CA, et al. Mutation of *mutL* homolog in hereditary colon cancer. *Science* 1994;263:1625-9.
 66. Nicolaides NC, Papadopoulos N, Liu B, Wei YF, Carter KC, Ruben SM, et al. Mutations of two *PMS* homologues in hereditary nonpolyposis colon cancer. *Nature* 1994;371:75-80.
 67. Eshleman JR, Lang EZ, Bowerfind GK, Parsons R, Vogelstein B, Willson JK, et al. Increased mutation rate at the *hprt* locus accompanies microsatellite instability in colon cancer. *Oncogene* 1995;10:33-7.
 68. Lanspa SJ, Jenkins JX, Cavalieri RJ, Smyrk TC, Watson P, Lynch J, et al. Surveillance in Lynch syndrome: how aggressive? *Am J Gastroenterol* 1994;89:1978-80.
 69. Aaltonen LA, Peltomäki P, Leach FS, Sistonen P, Pylkkanen L, Mecklin JP, et al. Clues to the pathogenesis of familial colorectal cancer. *Science* 1993;260:812-6.
 70. Aaltonen LA, Peltomäki P, Mecklin JP, Jarvinen H, Jass JR, Green JS, et al. Replication errors in benign and malignant tumors from hereditary nonpolyposis colorectal cancer patients. *Cancer Res* 54:1645-8.
 71. Ionov Y, Peinado MA, Malkhosyan S, Shibata D, Perucho M. Ubiquitous somatic mutations in simple repeated sequences reveal a new mechanism for colonic carcinogenesis. *Nature* 1993;363:558-61.
 72. Lothe RA, Peltomäki P, Meling GI, Aaltonen LA, Nystrom-Lahti M, Pylkkanen L, et al. Genomic instability in colorectal cancer: relationship of clinicopathological variables and family history. *Cancer Res* 1993;53:5849-52.
 73. Kim H, Jen J, Vogelstein B, Hamilton SR. Clinical and pathological characteristics of sporadic colorectal carcinomas with DNA replication errors in microsatellite sequences. *Am J Pathol* 1994;145:148-56.
 74. Thibodeau SN, Bren G, Schaid D. Microsatellite instability in cancer of the proximal colon. *Science* 1993;260:816-9.
 75. Liu B, Nicolaidis NC, Markowitz S, Wilson JK, Parsons RE, Jen J, et al. Mismatch repair gene defects in sporadic colorectal cancers with microsatellite instability. *Nat Genet* 1995;9:48-55.
 76. Bubb VJ, Curtis LJ, Cunningham C, Dunlop MG, Carothers AD, Morris RG, et al. Microsatellite instability and the role of *hMSH2* in sporadic colorectal cancer. *Oncogene* 1996;12:2641-9.
 77. Malkhosyan S, Rampino N, Yamamoto H, Perucho M. Frameshift mutator mutations. *Nature* 1996;382:499-500.
 78. Yamamoto H, Sawai H, Perucho M. Frameshift somatic mutations in gastrointestinal cancer of the microsatellite mutator phenotype. *Cancer Res* 1997;57:4420-6.
 79. Yamamoto H, Sawai H, Weber TK, Rodriguez-Bigas MA, Perucho M. Somatic frameshift mutations in DNA mismatch repair and proapoptosis genes in hereditary nonpolyposis colorectal cancer. *Cancer Res* 1998;58:997-1003.
 80. Kroser JA, Bachwich DR, Lichtenstein GR. Risk factors for the development of colorectal carcinoma and their modification. *Hematol Oncol Clin North Am* 1997;11:547-77.
 81. Horovitz IL, Huber M. Cancer of the colon: a study of age-related incidence. *Dis Colon Rectum* 1980;23:76-9.
 82. Boyle P, Zaridze DG. Colorectal cancer as a disease of the environment. *Ecol Dis* 1983;2:241-8.
 83. Whittemore AS, Wu-Williams AH, Lee M, Zheng S, Gallagher RP, Jiao DA, et al. Diet, physical activity, and colorectal cancer among Chinese in North America and China. *J Natl Cancer Inst* 1990;82:915-26.
 84. Bidoli E, Franceschi S, Talamini R, Barra S, La Vecchia C. Food consumption and cancer of the colon and rectum in north-eastern Italy. *Int J Cancer* 1992;50:223-9.
 85. Thun MJ, Calle EE, Namboodiri MM, Flanders WD, Coates RJ, Byers T, et al. Risk factors for fatal colon cancer in a large prospective study. *J Natl Cancer Inst* 1992;84:1491-500.
 86. Hague A, Manning AM, Hanlon KA, Huschtscha LI, Hart D, Paraskeva C. Sodium butyrate induces apoptosis in human colonic tumour cell lines in a p53-independent pathway: implications for the possible role of dietary fibre in the prevention of large-bowel cancer. *Int J Cancer* 1993;55:498-505.
 87. Jansson B. Geographic mappings of colorectal cancer rates: a retrospect of studies, 1974-1984. *Cancer Detect Prev* 1985;8:341-8.
 88. Nelson RL. Dietary minerals and colon carcinogenesis. *Anticancer Res* 1987;7:259-69.
 89. Nomura A, Heilbrun LK, Morris JS, Stemmermann GN. Serum selenium and the risk of can-

- cer, by specific sites: case-control analysis of prospective data. *J Natl Cancer Inst* 1987;79:103-8.
90. Folkman J. What is the evidence that tumors are angiogenesis dependent? *J Natl Cancer Inst* 1990;82:4-6.
 91. Graeber TG, Osmanian C, Jacks T, Housman DE, Koch CJ, Lowe SW, et al. Hypoxia-mediated selection of cells with diminished apoptotic potential in solid tumours. *Nature* 1996;379:88-91.
 92. Dameron KM, Volpert OV, Tainsky MA, Bouck N. Control of angiogenesis in fibroblasts by p53 regulation of thrombospondin-1. *Science* 1994;265:1582-4.
 93. Van Meir EG, Polverini PJ, Chazin VR, Su Huang HJ, de Tribolet N, Cavenee WK. Release of an inhibitor of angiogenesis upon induction of wild type p53 expression in glioblastoma cells. *Nat Genet* 1996;8:171-6.
 94. Holmgren L, O'Reilly MS, Folkman J. Dormancy of micrometastases: balanced proliferation and apoptosis in the presence of angiogenesis suppression. *Nat Med* 1995;1:149-53.
 95. Frank RE, Saclarides TJ, Leurgans S, Speziale NJ, Drab EA, Rubin DB. Tumor angiogenesis as a predictor of recurrence and survival in patients with node-negative colon cancer. *Ann Surg* 1995;222:695-9.
 96. Ebert EC, Brolin RE, Roberts AI. Characterization of activated lymphocytes in colon cancer. *Clin Immunol Immunopathol* 1989;50:72-81.
 97. Hakansson L, Adell G, Boeryd B, Sjogren F, Sjodahl R. Infiltration of mononuclear inflammatory cells into primary colorectal carcinomas: an immunohistological analysis. *Br J Cancer* 1997;75:374-80.
 98. Koch B, Giedl J, Hermanek P, Kalden JR. The analysis of mononuclear cell infiltrations in colorectal adenocarcinoma. *J Cancer Res Clin Oncol* 1985;109:42-51.
 99. Smith ME. MHC antigen expression in colorectal tumours. *Sem Cancer Biol* 1991;2:17-21.
 100. Giovannucci E, Egan KM, Hunter DJ, Stampfer MJ, Colditz GA, Willett WC, et al. Aspirin and the risk of colorectal cancer in women. *N Engl J Med* 1995;333:609-14.
 101. Giardiello FM, Hamilton SR, Krush AJ, Piantadosi S, Hyland LM, Celano P, et al. Treatment of colonic and rectal adenomas with sulindac in familial adenomatous polyposis. *N Engl J Med* 1993;328:1313-6.
 102. Eberhart CE, Coffey RJ, Radhika A, Giardiello FM, Ferrenbach S, DuBois RN. Up-regulation of cyclooxygenase 2 gene expression in human colorectal adenomas and adenocarcinomas. *Gastroenterology* 1994;107:1183-8.
 103. Pugh S, Thomas GA. Patients with adenomatous polyps and carcinomas have increased colonic mucosal prostaglandin E2. *Gut* 1994;35:675-8.
 104. Rigas B, Goldman IS, Levine L. Altered eicosanoid levels in human colon cancer. *J Lab Clin Med* 1993;122:518-23.
 105. Oshima M, Dinchuk JE, Kargman SL, Oshima H, Hancock B, Kwong E, et al. Suppression of intestinal polyposis in *APC*⁷¹⁶ knockout mice by inhibition of cyclooxygenase 2 (COX-2). *Cell* 1996;87:803-9.
 106. Kawamori T, Rao CV, Seibert K, Reddy BS. Chemopreventive activity of celecoxib, a specific cyclooxygenase-2 inhibitor, against colon carcinogenesis. *Cancer Res* 1998;58:409-12.
 107. Tsujii M, DuBois RN. Alterations in cellular adhesion and apoptosis in epithelial cells overexpressing prostaglandin endoperoxidase synthase 2. *Cell* 1995;83:493-501.
 108. Sheng H, Shao J, Morrow JD, Beauchamp RD, DuBois RN. Modulation of apoptosis and Bcl-2 expression by prostaglandin E2 in human colon cancer cells. *Cancer Res* 1998;58:362-6.
 109. Tsujii M, Kawano S, Tsuji S, Sawaoka H, Hori M, DuBois RN. Cyclooxygenase regulates angiogenesis induced by colon cancer cells. *Cell* 1998;93:705-16.
 110. Seed MP, Brown JR, Freemantle CN, Papworth JL, Colville-Nash PR, Willis D, et al. The inhibition of colon-26 adenocarcinoma development and angiogenesis by topical diclofenac in 2.5% hyaluronan. *Cancer Res* 1997;57:1625-9.

Review

Transforming growth factor- β and breast cancer Tumor promoting effects of transforming growth factor- β

Nancy Dumont and Carlos L Arteaga

Vanderbilt University School of Medicine and Vanderbilt-Ingram Cancer Center, Nashville, Tennessee, USA

Received: 3 December 1999

Revisions requested: 3 January 2000

Revisions received: 19 January 2000

Accepted: 26 January 2000

Published: 21 February 2000

Breast Cancer Res 2000, **2**:125–132

The electronic version of this article can be found online at
<http://breast-cancer-research.com/content/2/2/125>

© Current Science Ltd

Abstract

The transforming growth factor (TGF)- β s are potent growth inhibitors of normal epithelial cells. In established tumor cell systems, however, the preponderant experimental evidence suggests that TGF- β s can foster tumor-host interactions that indirectly support the viability and/or progression of cancer cells. The timing of this 'TGF- β switch' during the progressive transformation of epithelial cells is not clear. More recent evidence also suggests that autocrine TGF- β signaling is operative in some tumor cells, and can also contribute to tumor invasiveness and metastases independent of an effect on nontumor cells. The dissociation of antiproliferative and matrix associated effects of autocrine TGF- β signaling at a transcriptional level provides for a mechanism(s) by which cancer cells can selectively use this signaling pathway for tumor progression. Data in support of the cellular and molecular mechanisms by which TGF- β signaling can accelerate the natural history of tumors will be reviewed in this section.

Keywords: transforming growth factor (TGF)- β , TGF- β receptors, epithelial-to-mesenchymal transition, angiogenesis

Introduction

Although the transforming growth factor (TGF)- β s can be tumor suppressive [1], there is increasing evidence that TGF- β secretion by tumor cells and/or stromal cells within the peritumoral microenvironment can contribute to tumor maintenance and progression. How, then, can TGF- β s be both tumor suppressive and tumor promoting? This apparent paradox is reconciled by a study showing that, in a mouse skin model of chemical carcinogenesis, targeted expression of TGF- β 1 in suprabasal keratinocytes appears to have dual effects. It suppresses the formation of benign skin tumors, but once tumors develop, it enhances their progression to a highly invasive spindle cell phenotype [2*]. These results suggest that the effects of TGF- β 1 are

biphasic: TGF- β 1 acts early as a tumor suppressor, probably by inhibiting the proliferation of nontransformed cells, and it acts later as a tumor promoter by eliciting an epithelial-to-mesenchymal transition (EMT). Additional experiments have suggested that upregulation of TGF- β 3 in the spindle carcinomas was responsible for maintenance of this invasive phenotype [2*]. This is consistent with TGF- β 3 expression at sites in mouse embryos where epithelial-mesenchymal interactions are important, like the lung and palatal shelves [3,4], and also the abnormal lung development and cleft palate observed in TGF- β 3 null mice [5]. Also consistent with an early tumor suppressive effect is the recent observation that *tgf- β 1*^{-/-} mice develop an accelerated progression of epithelial hyperplasia to

CTL = cytotoxic T lymphocyte; EMT = epithelial-to-mesenchymal transition; JNK = c-Jun N-terminal kinase; MMP = matrix metalloproteases; PAI-1 = plasminogen activator inhibitor; PTHrP = parathyroid hormone-related protein; T β RI = TGF- β receptor type I; T β RII = TGF- β receptor type II; TGF- β = transforming growth factor- β .

colonic adenomas and cancers [6*]. The existence of dual effects for TGF- β s in tumor progression follows the observation that TGF- β -induced growth inhibitory responses and extracellular matrix responses may represent distinct processes in certain cell types. For example, overexpression of the antagonistic Smad, *Smad7*, in pancreatic carcinoma cell lines not only suppresses TGF- β 1-mediated growth inhibition, but enhances the ability of TGF- β 1 to induce matrix associated transcriptional responses [7*].

The progression of epithelial tumors to an invasive metastatic state is often associated with EMT, downregulation of cellular adhesion molecules, elevated expression of metalloproteases, and increased motility and angiogenesis, all of which can be modulated by TGF- β s. It is therefore not surprising that the TGF- β s can also promote tumorigenesis by modulating these critical processes. In support of this view, elevated levels of TGF- β are often observed in advanced carcinomas, and have been correlated with disease progression in several studies [8–13]. This suggests that secreting higher levels of TGF- β may provide an advantage to tumor cells. Both autocrine and paracrine signaling may be involved in conferring this selective advantage. While mutations in various components of the TGF- β signaling pathway have been observed in some carcinomas, particularly colorectal cancers [14,15], an intact TGF- β signaling pathway is often retained in other malignancies as some tumors can exhibit increased invasiveness in response to exogenous TGF- β [16,17*,18,19,20*,21]. Moreover, in a recent study of a large cohort of human breast tumors, loss or low levels of the type II TGF- β receptor (T β RII) correlated with high tumor grade, but 60% of *in situ* and invasive breast carcinomas retained robust levels of T β RII expression by immunohistochemistry [22]. Finally, although *Smad4* is frequently inactivated in pancreatic cancers [23,24], the *Smad* genes, which encode proteins that transduce TGF- β signals, are rarely mutated in most human carcinomas [25–30]. This suggests that after cells lose their sensitivity to TGF- β growth inhibition, autocrine TGF- β signaling may potentially promote tumor progression. In addition, TGF- β s produced in excess by tumor cells may act in a paracrine fashion on the peritumoral stroma, tumor neovessels, or the immune system, indirectly fostering tumor progression.

Autocrine effects

Epithelial-to-mesenchymal transition

Similar to keratinocytes [2*], TGF- β 1 can also induce a rapid and reversible EMT in melanoma cells [31], and in both nontumor [32] and Ha-Ras transformed [17*] mammary epithelial cells *in vitro*. In Ha-Ras mammary tumors, EMT appears to be initiated by TGF- β produced by peritumoral host cells and later maintained by autocrine TGF- β 1 as the converted tumor cells themselves begin to secrete TGF- β 1. The Ha-Ras tumor cells obtained after

EMT *in vitro* or *in vivo* display loss of epithelial polarity, downregulation of E-cadherin, disruption of cell–cell adhesion, and invasive properties in several *in vitro* assays [17*]. Supporting the importance of autocrine TGF- β for the tumorigenesis of Ha-Ras mammary cells, introduction of dominant negative T β RII into these cells retarded tumor formation and prevented EMT *in vivo*; moreover, introduction of the same construct into highly invasive murine colon carcinoma cells reconstituted an epithelial phenotype *in vitro*, and inhibited both tumor outgrowth and the establishment of metastases [20*]. In colon cancer cells of low invasive potential and with naturally occurring mutations in the T β RII gene, re-expression of T β RII function restored tumor cell invasiveness [20*]. In another study, expression of dominant negative T β RII in clones derived from a metastatic squamous carcinoma cell line prevented their spontaneous progression to a spindle phenotype *in vivo* [21]. Furthermore, approximately 90% of colon cancers with microsatellite instability have inactivating mutations of T β RII [33], and this instability is significantly correlated with longer patient survival [34], suggesting that complete loss of T β RII in carcinomas may limit systemic metastases. Taken together, these results suggest that EMT, local tumor growth, and metastatic progression can be sustained by autocrine TGF- β signaling.

When tumors are grown in nude mice, TGF- β s made by host cells can induce responses in tumor cells with intact TGF- β signaling, with the net effect of these tumor–host interactions being deleterious to the host. For example, MDA-231 human breast tumor cells secrete parathyroid hormone-related protein (PTHrP) in response to exogenous TGF- β 1, metastasize to bone when injected into nude mice, and induce osteolysis and hypercalcemia, resulting in host death. Transfection of these cells with dominant negative T β RII blocks TGF- β 1-mediated stimulation of PTHrP production. Mice injected with these cells exhibited less osteolysis, higher body weight, lower serum calcium and PTHrP levels, and longer survival than mice injected with control MDA-231 cells [35*]. On the contrary, accelerated osteolysis and reduced host survival were observed when mice were injected with tumor cells transfected with a constitutively active T β RI, suggesting a possible role for TGF- β -mediated responses in the pathogenesis of some adverse paraneoplastic syndromes.

Several recent studies have contributed to our understanding of the biochemical mechanisms by which transformed cells can lose autocrine growth inhibition but retain TGF- β -mediated responses that contribute to tumor progression. For example, oncogenic activation of the Ras pathway, acting via MAP kinases, causes phosphorylation of Smad2 and Smad3 at specific Erk consensus sites in the linker region between their DNA binding and transcriptional activation domains. This results in loss of nuclear accumulation of Smad2/3 and silencing of TGF- β -mediated

antiproliferative responses [36**]. In nontransformed mammary cells, introduction of mutant Ras not only blocks growth inhibition by TGF- β , but also subverts this pathway into one that can stimulate epithelial-to-mesenchymal transdifferentiation [17*,20*]. In MDCK epithelial cells, transfection of the missense mutations Smad2.D450E and Smad2.P445H, reported in primary colorectal and lung carcinomas, does not abolish TGF- β -mediated growth arrest. Instead, it increases both basal and TGF- β stimulated invasiveness, neither of which is prevented by overexpression of the inhibitory Smad7 [37*]. This suggests the existence of Smad 'gain-of-function' mutations that enhance malignant progression by mechanisms independent of T β RI and Smad phosphorylation. Another study has shown that Smad7 mRNA levels are increased in human pancreatic cancers compared with normal pancreas [7*]. Stable transfection of COLO-357 human pancreatic cancer cells with a Smad7 expression vector results in loss of TGF- β 1-mediated growth inhibition and p21/Cip1 promoter activity. However, TGF- β 1-induced plasminogen activator inhibitor-1 (PAI-1) promoter activity is maintained and, more importantly, basal PAI-1 promoter activity, PAI-1 mRNA levels, anchorage independent colony growth, and tumorigenicity in nude mice, are all increased in the Smad7 transfected clones [7*]. This result suggests another potential mechanism, the overexpression of Smad7, for the segregation between antiproliferative and matrix associated TGF- β responses. In addition, overexpression of Smad4 in colon carcinoma cells does not reconstitute TGF- β -mediated antiproliferative responses [38*,39], but inhibits cell adhesion and spreading, reduces the levels of urokinase plasminogen activator and PAI-1, and prolongs tumor latency [39], suggesting an additional function for Smad4 in restraining genes involved in peritumor proteolysis and invasion. This is further supported by reports of homozygous deletion of T β RI or homozygous missense mutations of T β RII [40,41], each coexisting with deletions of Smad4 in individual tumors. The coexistence of these mutations in the same tumors would not be expected if the function of these two gene products (T β RII and Smad4 or T β RI and Smad4) was limited to a single common signal transduction pathway. Taken together, these studies suggest, first, that the threshold for loss of TGF- β antimitogenic effects is lower than that required to lose responses associated with cell adhesion, invasion, and metastases; second, that not one but multiple biochemical mechanisms can contribute to the enhancement or unmasking of the tumor promoting effects of autocrine TGF- β ; and, third, that some of these mechanisms may be independent of Smad function or T β RI phosphorylation. The identification of Smad dependent and independent genes causally involved in these TGF- β -mediated tumor promoting effects requires further research. Of note, Hocevar *et al* [42*] recently reported c-Jun N-terminal kinase (JNK) dependent TGF- β -induced fibronectin expression in cell lines lacking the Smad4 gene or protein expression.

Increased motility

TGF- β can stimulate the motility of many cell types *in vitro* [43–45], therefore suggesting that TGF- β production *in vivo* may enhance migration of tumor cells and metastatic potential. Indeed, cyclosporine treatment of lung adenocarcinoma cells results in increased cell motility and anchorage independent growth *in vitro*, as well as increased metastases *in vivo*, all of which can be blocked with neutralizing TGF- β 1 antibodies [46]. These results suggest that *in vivo* tumor progression by cyclosporine is dependent on autocrine TGF- β 1. In prostate cancer cells, TGF- β 1 stimulates motility without affecting cell proliferation, suggesting that the effects on motility and proliferation may occur via different biochemical pathways [43].

Whether blockade of the Smad pathway, critical for TGF- β -mediated antimitogenic effects [47,48], is also critical for the effects of TGF- β s on cell motility is not clear. Some evidence suggests that the latter may follow alternative signaling pathways, perhaps in cooperation with activated oncogenes. Atfi *et al* [49] reported recently that inactivating components of the JNK pathway, which regulates AP-1 activity via c-Jun, inhibits TGF- β -mediated induction of 3TP-Lux, a reporter construct that contains Smad and AP-1 binding elements. Dominant negative mutants of RhoA, Rac1, and Cdc42, GTPases that mediate cell shape, cytoskeletal organization, and motility, abolish TGF- β -mediated transcription of AP-1 [49,50], suggesting that the Rho family of GTPases and the JNK pathway are essential components of TGF- β signaling responses. TGF- β 1 can also upregulate integrin linked kinase [31], a protein associated with fibronectin production and increased cell motility. In another study, TGF- β 1 treatment of NMuMG mouse mammary epithelial cells increased the expression of N-cadherin [51], which has been shown to increase motility of squamous cancer cells [52].

Paracrine effects

Induction of metalloproteases

Matrix metalloproteases (MMPs) play a critical role in the proteolytic degradation of basement membrane that is required for tumor invasion [53]. The expression of several MMPs, including MMP-2 [54] and MMP-9 [18,31,55], can be induced by TGF- β . Moreover, TGF- β 1 has been shown to selectively induce MMP-9 activity in a subset of metastatic but not primary mouse prostate tumors, implying that this TGF- β 1-induced response may be an important selection step in tumor progression [18]. There is also evidence that TGF- β increases MT-MMP-1 and MMP-9 expression in metastatic melanoma [31]. Although MMPs are listed separately, recent data implicate them strongly in the process of tumor-induced neovascularization [56], thereby suggesting that their upregulation might be an integral component of the TGF- β -mediated angiogenic processes discussed next.

Tumor angiogenesis

It is generally accepted that solid tumors require an adequate blood supply in order to grow beyond a few millimeters in size. TGF- β s, particularly TGF- β 1, have been shown to regulate new blood vessel formation both *in vitro* and *in vivo* by a combination of responses that include increased production and facilitation of vascular endothelial growth factor, facilitation of basic fibroblast growth factor mediated capillary sprouting, inhibition of endothelial cell migration, and increased production of extracellular matrix, among others (reviewed in [57]). In most cells, T β RI/ALK-5 is the signaling receptor for TGF- β . However, in endothelial cells, it has been suggested that ALK-1 may also function as a type I receptor for TGF- β [58]. In addition to the type I, II, and III TGF- β receptors, endoglin is another integral membrane protein that binds TGF- β 1 and TGF- β 3, and is highly expressed in endothelial cells [59]. Although TGF- β effects appear to be mediated mostly by the receptor specific Smad2 and Smad3 proteins [47,48], there is evidence that Smad5 is involved in TGF- β signaling in hematopoietic cells [60]. Targeted disruption of genes encoding various components of the TGF- β signaling pathway, including TGF- β 1 itself [61], its receptors, T β RII [62], ALK-1 [63], and endoglin [64], and one of its signal transducers, Smad5 [65], has each revealed that these proteins play an important role in vascular development. The phenotype of the TGF- β 1 and T β RII knockout mice is virtually indistinguishable and is characterized by defective endothelial differentiation resulting in abnormal capillary tube formation [61,62]. In contrast, disruption of ALK-1, endoglin, or Smad5 does not affect endothelial differentiation or vasculogenesis, but instead they each affect angiogenesis. In addition, *endoglin*^{-/-} and *Smad5*^{-/-} mice exhibit impaired vascular smooth muscle cell development. These results are consistent with previous reports demonstrating that TGF- β can regulate smooth muscle cell differentiation and migration *in vitro* [66], thus contributing to pericyte recruitment and vessel stabilization. This hypothesis, as it applies to tumor angiogenesis, is somewhat challenged by the notion that the majority of intratumoral neovessels seem to lack periendothelial smooth muscle cells [67], suggesting that there may be additional roles for the TGF- β s in tumor angiogenesis. In that light, Higaki and Shimokado [68] recently reported TGF- β 1-mediated stimulation of phosphatidylinositol-3 kinase activity and amino acid uptake in vascular smooth muscle cells, suggesting a direct anti-apoptotic role for TGF- β . Elucidation of the paracrine mechanisms driving TGF- β -mediated tumor angiogenesis requires further investigation.

Further supporting the role of TGF- β s in tumor angiogenesis, administration of a neutralizing TGF- β 1 antibody to nude mice harboring CHO cell xenografts transfected with ectopic TGF- β 1 inhibits both tumor growth and intratumor microvessel density [69]. In addition, a monoclonal anti-

body that blocks TGF- β 1, TGF- β 2, and TGF- β 3 has been shown to suppress the growth of TGF- β 1-overexpressing renal cancer xenografts [70]. In this study, the TGF- β blocking monoclonal-abrogated factor VIII staining in the xenografts, suggesting an antitumor mechanism that targets endothelial cells [70]. Furthermore, TGF- β 1 and PAI-1 have been shown to inhibit the conversion of plasminogen to the anti-angiogenic molecule angiostatin in medium conditioned by human pancreatic cancer cells [71]. This suggests an additional pro-angiogenic mechanism for TGF- β by interfering with the production of endogenous inhibitors of endothelial cell proliferation. Finally, high levels of TGF- β 1 mRNA correlate strongly with high microvessel density in breast tumors, and each of these factors is associated with poor patient outcome [72].

Host immunosuppression

TGF- β 1 and TGF- β 2 are potent immunosuppressants [73]. Thus, elevated levels of TGF- β s secreted by tumors could potentially inhibit immune effector cells and favor tumor progression. In support of this idea, Torre Amione *et al* [74] demonstrated that, unlike parental tumor cells, fibrosarcoma cells transfected to express 10 ng/ml TGF- β 1 *in vitro* are unable to induce cytotoxic T lymphocyte (CTL) responses and can escape immune recognition. Likewise, EMT6 mammary tumor cells, which produce high levels of TGF- β 1, can inhibit CTLs *in vivo*. Transfection of these cells with interleukin-2, a known T cell growth factor, can reverse this TGF- β 1 effect and induce tumor rejection [75]. This result suggests that, by dampening the generation of tumor reactive T cells, TGF- β can promote tumor viability. There is also evidence that overexpression of the soluble T β RII extracellular domain in thymoma cells can prevent the progression of unmodified thymoma cells when injected near the primary tumor inoculation site [76], further suggesting that secretion of soluble T β RII by these cells is sufficient to restore tumor specific cellular immunity and mediate partial tumor rejection. Overall, these results are consistent with the phenotype of *TGF- β 1* null mice that die shortly after birth as a result of widespread inflammation and multiorgan T cell infiltration and necrosis [77].

In addition to inhibiting CTL responses, TGF- β s can modulate other immune functions that may favor tumor progression. For example, CHO cells transfected with an expression vector encoding latent TGF- β 1, when injected into nude mice, can decrease mouse spleen natural killer activity and rapidly form tumors [78]. Antagonizing TGF- β s by intraperitoneal injection of an antibody that neutralizes TGF- β 1, TGF- β 2, and TGF- β 3 has the opposite effect. It prevents tumor and metastases formation by MDA-231 human breast carcinoma cells, and markedly increases natural killer activity of mouse splenocytes [79]. Consistent with this TGF- β -mediated immunosuppressive effect, reduced immune function has been observed in animals

bearing TGF- β overexpressing tumors [80] as well as in patients with glioblastoma, a common type of brain tumor that frequently overexpresses TGF- β 2 [81].

The cited studies suggest that tumor cell secreted TGF- β s may block the efferent function of immune effectors at sites of tumor implantation. Other reports, however, suggest tumor cell TGF- β s may modify the afferent component of the immune response and confer antitumor immunity. Stable infection of breast and glioma tumor cells with antisense TGF- β 1 and antisense TGF- β 2 retroviruses, respectively, has been shown to restore the immunogenicity of these tumor cells when injected into immunocompetent animals. Furthermore, they induce a partial rejection of unmodified, less immunogenic established wild type tumor cells [82,83]. In both of these studies, *in vitro* and *in vivo* CTL activity was markedly increased in medium conditioned by antisense TGF- β -infected cells and/or in mice injected with tumor cells bearing the antisense compared with tumor cells infected with a control vector. These studies have therapeutic implications for the use of an antisense TGF- β based approach as a means of adoptive immunotherapy against TGF- β overproducing tumors.

Alternative views and conclusions

A tumor permissive role for the TGF- β s may not apply to all solid tumors. Indeed, transfection of an antisense TGF- β 1 expression vector into FET and CBS well-differentiated human colon cancer cells has been shown to enhance tumor formation in nude mice [84,85], supporting the notion that, in some fully transformed cells, endogenous TGF- β 1 can continue to mediate a tumor suppressor function. In a recent report, mice bearing transplanted gallbladder Mz-Cha-2 tumors showed inhibition of angiogenesis and leukocyte-endothelial cell interactions at a distant cranial site and threefold higher levels of circulating TGF- β 1 compared with tumor free mice [86]. This reduction in microvessel density and leukocyte rolling were reversed by systemic administration of a TGF- β 1 neutralizing antibody, suggesting a negative role for TGF- β 1 in early neovascularization. Moreover, in a recent survey of 104 *in situ* and invasive primary breast carcinomas, 40/45 (89%) tumors with low invasive potential and low proliferation rate exhibited high levels of T β RII by immunohistochemistry [22]. Whether autocrine TGF- β signaling is causally associated with the observed low proliferation and invasiveness in this subset of breast tumors is a question that remains unclear.

Nonetheless, the potential tumor promoting effects of TGF- β provide novel molecular targets for interventions aimed at altering the natural history of solid tumors. The lack of an obvious physiological role for TGF- β signaling in postdevelopmental normal physiological states suggests that these interventions may in fact be tumor specific and

spare the tumor host from undue toxicity. Several approaches have been proposed, including the use of blocking antibodies against TGF- β 1, TGF- β 2, and TGF- β 3, using the soluble ectodomains of the type II and III TGF- β receptors, which would sequester TGF- β isoforms at tumor sites and prevent binding to cognate receptors [87,88], and, finally, using adenovirus encoding inhibitors of TGF- β signaling [89], to name a few. The theoretical and logistical strengths and limitations of these approaches are beyond the scope of this review. Nonetheless, these represent tools that, if effective in blocking TGF- β action, will allow us to address the net effect of autocrine/paracrine TGF- β signaling at early and late stages of transformation and cancer progression.

References

Articles of particular interest have been highlighted as:

- of special interest
- of outstanding interest

1. Markowitz SD, Roberts AB: **Tumor suppressor activity of the TGF-beta pathway in human cancers.** *Cytokine Growth Factor Rev* 1996, 7:93-102.
2. Cui W, Fowlis DJ, Bryson S, *et al*: **TGFbeta1 inhibits the formation of benign skin tumors, but enhances progression to invasive spindle carcinomas in transgenic mice.** *Cell* 1996, 86:531-542.
This is the first study which demonstrates that TGF- β 1 may actually have a dual role and be both tumor suppressive and tumor promoting. Mice expressing active TGF- β 1 under the control of the K1V promoter, specific to suprabasal skin keratinocytes, are more resistant to the induction of benign skin tumors than controls. However, their conversion to invasive, spindle cell carcinomas is markedly increased.
3. Schmid P, Cox D, Bilbe G, Maier R, McMaster GK: **Differential expression of TGF beta 1, beta 2 and beta 3 genes during mouse embryogenesis.** *Development* 1991, 111:117-130.
4. Millan FA, Denhez F, Kondaiah P, Akhurst RJ: **Embryonic gene expression patterns of TGF beta 1, beta 2 and beta 3 suggest different developmental functions in vivo.** *Development* 1991, 111:131-143.
5. Kaartinen V, Voncken JW, Shuler C, *et al*: **Abnormal lung development and cleft palate in mice lacking TGF-beta 3 indicates defects of epithelial-mesenchymal interaction.** *Nat Genet* 1995, 11:415-421.
6. Engle SJ, Hoving JB, Boivin GP, Ormsby I, Gartside PS, Doetschman T: **Transforming growth factor beta1 suppresses nonmetastatic colon cancer at an early stage of tumorigenesis.** *Cancer Res* 1999, 59:3379-3386.
Tgfb1^{-/-} *Rag2*^{-/-} mice rapidly develop carcinomas of the cecum and colon. In contrast, the colon from *Tgfb*^{+/+} and *Tgfb*^{+/+} *Rag*^{-/-} mice remains hyperplastic with almost no progression to carcinoma. Data in the paper suggest that loss of TGF- β 1 contributes to tumor formation by mechanisms other than loss of growth control, increased inflammation, or genetic instability.
7. Kleeff J, Ishiwata T, Maruyama H, *et al*: **The TGF-beta signaling inhibitor smad7 enhances tumorigenicity in pancreatic cancer.** *Oncogene* 1999, 18:5363-5372.
Stable transfection of ectopic Smad7 in COLO-357 human pancreatic cancer cells abrogates TGF- β -mediated growth inhibition and transcription via the p21 promoter, but potentiates PAI-1 mRNA induction, 3TP-luciferase activity, and tumorigenicity in nude mice, highlighting a dissociation between the growth and matrix effects of TGF- β .
8. Walker RA, Dearing SJ: **Transforming growth factor beta 1 in ductal carcinoma in situ and invasive carcinomas of the breast.** *Eur J Cancer* 1992, 28:641-644.

9. Gorsch SM, Memoli VA, Stukel TA, Gold LI, Arrick BA: Immunohistochemical staining for transforming growth factor beta 1 associates with disease progression in human breast cancer. *Cancer Res* 1992, 52:6949-6952.
 10. Friess H, Yamanaka Y, Buchler M, et al: Enhanced expression of transforming growth factor beta isoforms in pancreatic cancer correlates with decreased survival. *Gastroenterology* 1993, 105: 1846-1856.
 11. Reed JA, McNutt NS, Prieto VG, Albino AP: Expression of transforming growth factor beta 2 in malignant melanoma correlates with the depth of tumor invasion. Implications for tumor progression. *Am J Pathol* 1994, 145:97-104.
 12. Ito N, Kawata S, Tamura S, et al: Positive correlation of plasma transforming growth factor-beta 1 levels with tumor vascularity in hepatocellular carcinoma. *Cancer Lett* 1995, 89:45-48.
 13. Wikstrom P, Stattin P, Franck-Lissbrant I, Damber JE, Bergh A: Transforming growth factor beta1 is associated with angiogenesis, metastasis, and poor clinical outcome in prostate cancer. *Prostate* 1998, 37:19-29.
 14. Hata A, Shi Y, Massague J: TGF-beta signaling and cancer: structural and functional consequences of mutations in Smads. *Mol Med Today* 1998, 4:257-262.
 15. Dumont N: Genetic and epigenetic contributions to colorectal cancer. *APMIS* 1999, 107:711-722.
 16. Welch DR, Fabra A, Nakajima M: Transforming growth factor beta stimulates mammary adenocarcinoma cell invasion and metastatic potential. *Proc Natl Acad Sci USA* 1990, 87:7678-7682.
 17. Oft M, Peli J, Rudaz C, Schwarz H, Beug H, Reichmann E: TGF-beta1 and Ha-Ras collaborate in modulating the phenotypic plasticity and invasiveness of epithelial tumor cells. *Genes Dev* 1996, 10: 2462-2477.
- Ras-transformed mouse mammary epithelial cells lose the growth inhibitory response to TGF- β and undergo a TGF- β -induced EMT that is maintained by the upregulated expression of TGF- β in an autocrine fashion.
18. Sehgal I, Baley PA, Thompson TC: Transforming growth factor beta1 stimulates contrasting responses in metastatic versus primary mouse prostate cancer-derived cell lines *in vitro*. *Cancer Res* 1996, 56:3359-3365.
 19. Farina AR, Coppa A, Tiberio A, et al: Transforming growth factor-beta1 enhances the invasiveness of human MDA-MB-231 breast cancer cells by up-regulating urokinase activity. *Int J Cancer* 1998, 75:721-730.
 20. Oft M, Heider KH, Beug H: TGFbeta signaling is necessary for carcinoma cell invasiveness and metastasis. *Curr Biol* 1998, 8: 1243-1252.
- Inhibition of autocrine TGF- β signaling, by expression of dominant negative T β RII constructs, can revert the metastatic, invasive phenotype of tumor cells *in vitro* and *in vivo*, thus implying that the contribution of TGF- β signaling to invasion is a cell autonomous process.
21. Portella G, Cumming SA, Liddell J, et al: Transforming growth factor beta is essential for spindle cell conversion of mouse skin carcinoma *in vivo*: Implications for tumor invasion. *Cell Growth Diff* 1998, 9:393-404.
 22. Gobbi H, Arteaga CL, Jenson RA, et al: Loss of expression of transforming growth factor beta type II receptor correlates with high tumor grade in human breast *in situ* and invasive carcinomas. *Histopathology* 2000, in press.
 23. Hahn SA, Schutte M, Hoque AT, et al: DPC4, a candidate tumor suppressor gene at human chromosome 18q21.1. *Science* 1996, 271:350-353.
 24. Jonson T, Gorunova L, Dawiskiba S, et al: Molecular analyses of the 15q and 18q SMAD genes in pancreatic cancer. *Genes Chromosomes Cancer* 1999, 24:62-71.
 25. Kim SK, Fan Y, Papadimitrakopoulou V, et al: DPC4, a candidate tumor suppressor gene, is altered infrequently in head and neck squamous cell carcinoma. *Cancer Res* 1996, 56:2519-2521.
 26. Lei J, Zou TT, Shi YQ, et al: Infrequent DPC4 gene mutation in esophageal cancer, gastric cancer and ulcerative colitis-associated neoplasms. *Oncogene* 1996, 13:2459-2462.
 27. Schutte M, Hruban RH, Hedrick L, et al: DPC4 gene in various tumor types. *Cancer Res* 1996, 56:2527-2530.
 28. Riggins GJ, Kinzler KW, Vogelstein B, Thiagalingam S: Frequency of Smad gene mutations in human cancers. *Cancer Res* 1997, 57:2578-2580.
 29. Arai T, Akiyama Y, Okabe S, Ando M, Endo M, Yuasa Y: Genomic structure of the human Smad3 gene and its infrequent alterations in colorectal cancers. *Cancer Lett* 1998, 122:157-163.
 30. Latil A, Pesche S, Valeri A, Fournier G, Cussenot O, Lidereau R: Expression and mutational analysis of the MADR2/Smad2 gene in human prostate cancer. *Prostate* 1999, 40:225-231.
 31. Janji B, Melchior C, Gouon V, Vallar L, Kieffer N: Autocrine TGF-beta-regulated expression of adhesion receptors and integrin-linked kinase in HT-144 melanoma cells correlates with their metastatic phenotype. *Int J Cancer* 1999, 83:255-262.
 32. Miettinen PJ, Ebner R, Lopez AR, Derynck R: TGF-beta induced transdifferentiation of mammary epithelial cells to mesenchymal cells: involvement of type I receptors. *J Cell Biol* 1994, 127:2021-2036.
 33. Parsons R, Myeroff LL, Liu B, et al: Microsatellite instability and mutations of the transforming growth factor beta type II receptor gene in colorectal cancer. *Cancer Res* 1995, 55:5548-5550.
 34. Thibodeau SN, Bren G, Schaid D: Microsatellite instability in cancer of the proximal colon. *Science* 1993, 260:816-819.
 35. Yin JJ, Selander K, Chirgwin JM, et al: TGF-beta signaling blockade inhibits PTHrP secretion by breast cancer cells and bone metastases development. *J Clin Invest* 1999, 103:197-206.
- This paper reports a lethal bone metastases model of MDA-231 human breast tumor cells in athymic mice. Expression of ectopic dominant negative T β RII into MDA-231 cells inhibits TGF- β -induced PTHrP production by tumor cells, tumor cell mediated osteolysis, and prolongs mouse survival. On the contrary, forced expression of a constitutively active T β RI expression vector accelerates these processes.
36. Kretzschmar M, Doody J, Timokhina I, Massague J: A mechanism of repression of TGFbeta/Smad signaling by oncogenic Ras. *Genes Dev* 1999, 13:804-816.
- Ras, acting via Erk MAP kinases, causes phosphorylation at sites in the linker region of Smad2 and Smad3, which, in turn, inhibit Smad accumulation in the nucleus and TGF- β -mediated growth inhibition. Expression of a Ras-resistant Smad3 mutant, lacking the MAPK phosphorylation sites, restored transcriptional and cellular responses to exogenous TGF- β in Ras-transformed cells.
37. Prunier C, Mazars A, Noe V, et al: Evidence that Smad2 is a tumor suppressor implicated in the control of cellular invasion. *J Biol Chem* 1999, 274:22919-22922.
- The tumor-derived missense mutation Smad2.D450E is not a 'loss of function mutation'; that is, it does not confer resistance to TGF- β -mediated growth inhibition. Rather unexpectedly, it behaves as a 'gain of function' mutation by promoting increased tumor cell invasiveness. Overexpression of Smad7 does not block this effect on invasion, suggesting that Smad2.D450E action does not require T β RI phosphorylation.
38. Calonge MJ, Massague J: Smad4/DPC4 silencing and hyperactive ras jointly disrupt transforming growth factor-beta antiproliferative responses in colon cancer cells. *J Biol Chem* 1999, 274:33637-33643.
- Forced expression of Smad4 does not rescue TGF- β -mediated antiproliferative and transcriptional responses in SW480.7 colon cancer cells, which contain an activated Ki-ras oncogene. Cotransfection of Smad4 with a Ras

phosphorylation-resistant Smad3, however, restores TGF- β actions, suggesting that both loss of Smad4 function plus hyperactive Ras signaling partner in abrogating TGF- β antiproliferative responses.

39. Schwarte-Waldhoff I, Klein S, Blass-Kampmann S, et al: **DPC4/SMAD4 mediated tumor suppression of colon carcinoma cells is associated with reduced urokinase expression.** *Oncogene* 1999, **18**:3152-3158.
40. Goggins M, Shekher M, Turnacioglu K, Yeo CJ, Hruban RH, Kern SE: **Genetic alterations of the transforming growth factor beta receptor genes in pancreatic and biliary adenocarcinomas.** *Cancer Res* 1998, **58**:5329-5332.
41. Grady WM, Myeroff LL, Swinler SE, et al: **Mutational inactivation of transforming growth factor beta receptor type II in microsatellite stable colon cancers.** *Cancer Res* 1999, **59**:320-324.
42. Hoyer BA, Brown TL, Howe PH: **TGF-beta induces fibronectin synthesis through a c-Jun N-terminal kinase-dependent, Smad4 independent pathway.** *EMBO J* 1999, **18**:1345-1356.
First report of a TGF- β -regulated gene independent of Smad4 function.
43. Morton DM, Barrack ER: **Modulation of transforming growth factor beta 1 effects on prostate cancer cell proliferation by growth factors and extracellular matrix.** *Cancer Res* 1995, **55**:2596-2602.
44. Hasina R, Matsumoto K, Matsumoto-Taniura N, Kato I, Sakuda M, Nakamura T: **Autocrine and paracrine motility factors and their involvement in invasiveness in a human oral carcinoma cell line.** *Br J Cancer* 1999, **80**:1708-1717.
45. Zicha D, Genot E, Dunn GA, Kramer IM: **TGFbeta1 induces a cell-cycle-dependent increase in motility of epithelial cells.** *J Cell Sci* 1999, **112**:447-454.
46. Hojo M, Morimoto T, Maluccio M, et al: **Cyclosporine induces cancer progression by a cell-autonomous mechanism.** *Nature* 1999, **397**:530-534.
47. Liu X, Sun Y, Constantinescu SN, Karam E, Weinberg RA, Lodish HF: **Transforming growth factor beta-induced phosphorylation of Smad3 is required for growth inhibition and transcriptional induction in epithelial cells.** *Proc Natl Acad Sci USA* 1997, **94**:10669-10674.
48. Datto MB, Frederick JP, Pan L, Borton AJ, Zhuang Y, Wang XF: **Targeted disruption of Smad3 reveals an essential role in transforming growth factor beta-mediated signal transduction.** *Mol Cell Biol* 1999, **19**:2495-2504.
49. Atfi A, Djelloul S, Chastre E, Davis R, Gespach C: **Evidence for a role of Rho-like GTPases and stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK) in transforming growth factor beta-mediated signaling.** *J Biol Chem* 1997, **272**:1429-1432.
50. Engel ME, Datta PK, Moses HL: **RhoB is stabilized by transforming growth factor beta and antagonizes transcriptional activation.** *J Biol Chem* 1998, **273**:9921-9926.
51. Knudsen KA, Frankowski C, Johnson KR, Wheelock MJ: **A role for cadherins in cellular signaling and differentiation.** *J Cell Biochem Suppl* 1998, **30-31**:168-176.
52. Islam S, Carey TE, Wolf GT, Wheelock MJ, Johnson KR: **Expression of N-cadherin by human squamous carcinoma cells induces a scattered fibroblastic phenotype with disrupted cell-cell adhesion.** *J Cell Biol* 1996, **135**:1643-1654.
53. Liotta LA, Stetler-Stevenson WG: **Tumor invasion and metastasis: an imbalance of positive and negative regulation.** *Cancer Res* 1991, **51**(suppl):S5054-S5059.
54. Overall CM, Wrana JL, Sodek J: **Transcriptional and post-transcriptional regulation of 72-kDa gelatinase/type IV collagenase by transforming growth factor-beta 1 in human fibroblasts. Comparisons with collagenase and tissue inhibitor of matrix metalloproteinase gene expression.** *J Biol Chem* 1991, **266**:14064-14071.
55. Samuel SK, Hurta RA, Kondaiah P, et al: **Autocrine induction of tumor protease production and invasion by a metallothionein-regulated TGF-beta 1 (Ser223, 225).** *EMBO J* 1992, **11**:1599-1605.
56. Coussens LM, Raymond WW, Bergers G, et al: **Inflammatory mast cells up-regulate angiogenesis during squamous epithelial carcinogenesis.** *Genes Dev* 1999, **13**:1382-1397.
57. Pepper MS: **Transforming growth factor-beta: vasculogenesis, angiogenesis, and vessel wall integrity.** *Cytokine Growth Factor Rev* 1997, **8**:21-43.
58. Lux A, Attisano L, Marchuk DA: **Assignment of transforming growth factor beta1 and beta3 and a third new ligand to the type I receptor ALK-1.** *J Biol Chem* 1999, **274**:9984-9992.
59. Barbara NP, Wrana JL, Letarte M: **Endoglin is an accessory protein that interacts with the signaling receptor complex of multiple members of the transforming growth factor-beta superfamily.** *J Biol Chem* 1999, **274**:584-594.
60. Bruno E, Horrigan SK, Van Den Berg D, et al: **The Smad5 gene is involved in the intracellular signaling pathways that mediate the inhibitory effects of transforming growth factor-beta on human hematopoiesis.** *Blood* 1998, **91**:1917-1923.
61. Dickson MC, Martin JS, Cousins FM, Kulkarni AB, Karlsson S, Akhurst RJ: **Defective haematopoiesis and vasculogenesis in transforming growth factor-beta 1 knock out mice.** *Development* 1995, **121**:1845-1854.
62. Oshima M, Oshima H, Taketo MM: **TGF-beta receptor type II deficiency results in defects of yolk sac hematopoiesis and vasculogenesis.** *Dev Biol* 1996, **179**:297-302.
63. Oh SP, Seki T, Goss KA, et al: **ALK-1 modulates TGF-beta1 signaling during vascular development.** In *Proceedings of the Third International TGF-Beta Conference: Biological Mechanisms and Clinical Applications*. September 20-22 1999, NIH, Bethesda, MD.
64. Li DY, Sorensen LK, Brooke BS, et al: **Defective angiogenesis in mice lacking endoglin.** *Science* 1999, **284**:1534-1537.
65. Yang X, Castilla LH, Xu X, et al: **Angiogenesis defects and mesenchymal apoptosis in mice lacking SMAD5.** *Development* 1999, **126**:1571-1580.
66. Hirschi KK, Rohovsky SA, D'Amore PA: **PDGF, TGF-beta, and heterotypic cell-cell interactions mediate endothelial cell-induced recruitment of 10T1/2 cells and their differentiation to a smooth muscle fate [published erratum appears in J Cell Biol 1998, 141(5):1287].** *J Cell Biol* 1998, **141**:805-814.
In response to activated TGF- β 1, 10T1/2 fibroblasts migrate toward endothelial cells and acquire a smooth muscle phenotype. This effect is blocked by TGF- β neutralizing antibodies, suggesting that TGF- β s are critical for heterotypic cell-cell interactions involved in pericyte recruitment to nascent capillaries and in microvessel wall integrity.
67. Benjamin LE, Golijanin D, Itin A, Podes D, Keshet E: **Selective ablation of immature blood vessels in established human tumors follows vascular endothelial growth factor withdrawal.** *J Clin Invest* 1999, **103**:159-165.
68. Higaki M, Shimokado K: **Phosphatidylinositol 3-kinase is required for growth factor induced amino acid uptake by vascular smooth muscle cells.** *Arterioscler Thromb Vasc Biol* 1999, **19**:2127-2132.
69. Ueki N, Nakazato M, Ohkawa T, et al: **Excessive production of transforming growth factor beta 1 can play an important role in the development of tumorigenesis by its action for angiogenesis: validity of neutralizing antibodies to block tumor growth.** *Biochim Biophys Acta* 1992, **1137**:189-196.
70. Ananth S, Knebelmann B, Gruning W, et al: **Transforming growth factor beta1 is a target for the von Hippel-Lindau tumor suppressor and a critical growth factor for clear cell renal carcinoma.** *Cancer Res* 1999, **59**:2210-2216.

71. O'Mahony CA, Albo D, Tuszynski GP, Berger DH: Transforming growth factor-beta 1 inhibits generation of angiostatin by human pancreatic cancer cells. *Surgery* 1998, 124:388-393.
72. Relf M, LeJeune S, Scott PA, et al: Expression of the angiogenic factors vascular endothelial cell growth factor, acidic and basic fibroblast growth factor, tumor growth factor beta-1, platelet-derived endothelial cell growth factor, placenta growth factor, and pleiotrophin in human primary breast cancer and its relation to angiogenesis. *Cancer Res* 1997, 57:963-969.
73. Letterio JJ, Roberts AB: Regulation of immune responses by TGF-beta. *Annu Rev Immunol* 1998, 16:137-161.
74. Torre-Amione G, Beauchamp RD, Koeppen H, et al: A highly immunogenic tumor transfected with a murine transforming growth factor type beta 1 cDNA escapes immune surveillance. *Proc Natl Acad Sci USA* 1990, 87:1486-1490.
75. McAdam AJ, Felcher A, Woods ML, et al: Transfection of transforming growth factor-beta producing tumor EMT6 with interleukin-2 elicits tumor rejection and tumor reactive cytotoxic T-lymphocytes. *J Immunother Emphasis Tumor Immunol* 1994, 15:155-164.
76. Won J, Kim H, Park EJ, Hong Y, Kim SJ, Yun Y: Tumorigenicity of mouse thymoma is suppressed by soluble type II transforming growth factor beta receptor therapy. *Cancer Res* 1999, 59:1273-1277.
77. Shull MM, Ormsby I, Kier AB, et al: Targeted disruption of the mouse transforming growth factor-beta 1 gene results in multifocal inflammatory disease. *Nature* 1992, 359:693-699.
78. Wallick SC, Figari IS, Morris RE, Levinson AD, Palladino MA: Immunoregulatory role of transforming growth factor beta (TGF-beta) in development of killer cells: comparison of active and latent TGF-beta 1. *J Exp Med* 1990, 172:1777-1784.
79. Arteaga CL, Hurd SD, Winnier AR, Johnson MD, Fendly BM, Forbes JT: Anti-transforming growth factor (TGF)-beta antibodies inhibit breast cancer cell tumorigenicity and increase mouse spleen natural killer cell activity. Implications for a possible role of tumor cell/host TGF-beta interactions in human breast cancer progression. *J Clin Invest* 1993, 92:2569-2576.
80. Li XF, Takiuchi H, Zou JP, et al: Transforming growth factor-beta (TGF-beta) mediated immunosuppression in the tumor-bearing state: enhanced production of TGF-beta and a progressive increase in TGF-beta susceptibility of anti-tumor CD4⁺ T cell function. *Jpn J Cancer Res* 1993, 84:315-325.
81. Kuppner MC, Hamou MF, Bodmer S, Fontana A, de Tribolet N: The glioblastoma derived T-cell suppressor factor/transforming growth factor beta 2 inhibits the generation of lymphokine-activated killer (LAK) cells. *Int J Cancer* 1988, 42:562-567.
82. Fakhrai H, Dorigo O, Shawler DL, et al: Eradication of established intracranial rat gliomas by transforming growth factor beta antisense gene therapy. *Proc Natl Acad Sci USA* 1996, 93:2909-2914.
83. Park JA, Wang E, Kurt RA, Schluter SF, Hersch EM, Akporiaye ET: Expression of an antisense transforming growth factor-beta1 transgene reduces tumorigenicity of EMT6 mammary tumor cells. *Cancer Gene Ther* 1997, 4:42-50.
84. Wu SP, Theodorescu D, Kerbel RS, et al: TGF-beta 1 is an autocrine-negative growth regulator of human colon carcinoma FET cells *in vivo* as revealed by transfection of an antisense expression vector. *J Cell Biol* 1992, 116:187-196.
85. Wu SP, Sun LZ, Willson JK, Humphrey L, Kerbel R, Brattain MG: Repression of autocrine transforming growth factor beta 1 and beta 2 in quiescent CBS colon carcinoma cells leads to progression of tumorigenic properties. *Cell Growth Diff* 1993, 4:115-123.
86. Gohongi T, Fukumura D, Boucher Y, et al: Tumor-host interactions in the gallbladder suppress distal angiogenesis and tumor growth: involvement of transforming growth factor beta1. *Nat Med* 1999, 5:1203-1208.
87. Bandyopadhyay A, Zhu Y, Cibull ML, Bao L, Chen C, Sun L: A soluble transforming growth factor beta type III receptor suppresses tumorigenicity and metastasis of human breast cancer MDA-MB-231 cells. *Cancer Res* 1999, 59:5041-5046.
88. George J, Roulot D, Kotliansky VE, Bissell DM: *In vivo* inhibition of rat stellate cell activation by soluble transforming growth factor beta type II receptor: a potential new therapy for hepatic fibrosis. *Proc Natl Acad Sci USA* 1999, 96:12719-12724.
89. Nakao A, Fujii M, Matsumura R, et al: Transient gene transfer and expression of Smad7 prevents bleomycin-induced lung fibrosis in mice. *J Clin Invest* 1999, 104:5-11.

Authors' affiliations: Nancy Dumont (Department of Cell Biology, Vanderbilt University School of Medicine, Nashville, Tennessee, USA) and Carlos L Arteaga (Departments of Cell Biology and Medicine, Vanderbilt University School of Medicine, Department of Veteran Affairs Medical Center, and Vanderbilt-Ingram Cancer Center, Nashville, Tennessee, USA)

Sponsorship: This review was supported by US Army DAMD17-98-1-8263 Pre-doctoral Training Award (to ND), PHS grant R01 CA62212, US Army DAMD17-98-1-8262 IDEA grant, a Department of Veteran Affairs Clinical Investigator Award (to CLA), and a Vanderbilt-Ingram Cancer Center support grant CA68485

Correspondence: Carlos L Arteaga, Division of Hematology-Oncology/VUMC, 22nd Avenue South, 1956 TVC, Nashville, TN 37232-5536, USA. Tel: +1 615 936 3524/1919; fax: +1 615 936 1790; e-mail: carlos.arteaga@mcmail.vanderbilt.edu

INVASION AND METASTASIS OF A MAMMARY TUMOR INVOLVES TGF- β SIGNALING

Julie A. McEARCHER¹, James J. KOBIE¹, Vivian MACK¹, Rita S. WU¹, Linda MEADE-TOLLIN², Carlos L. ARTEAGA⁴, Nancy DUMONT⁴, David BESSELS³, Elisabeth SEFTOR⁵, Mary J.C. HENDRIX⁵, Emmanuel KATSANIS⁶ and Emmanuel T. AKPORIAYE^{1*}

¹Department of Microbiology and Immunology, University of Arizona, Tucson, AZ

²Department of Surgery, University of Arizona, Tucson, AZ

³Department of Veterinary Sciences and Microbiology and University Animal Care, University of Arizona, Tucson, AZ

⁴Departments of Medicine and Cell Biology, Vanderbilt University School of Medicine, Department of Veteran Affairs Medical Center and Vanderbilt-Ingram Cancer Center (C.L.A.), Nashville, TN

⁵Department of Anatomy and Cell Biology, University of Iowa, Iowa City, IA

⁶Department of Pediatrics, University of Arizona, Tucson, AZ

Several studies have correlated escape from TGF- β -mediated cell cycle arrest with the tumorigenic phenotype. Most often, this escape from growth control has been linked to dysfunctional TGF- β receptors or defects in the TGF- β -mediated SMAD signaling pathway. In this report, we found that highly metastatic 4T1 mammary carcinoma cells express functional TGF- β receptors capable of initiating SMAD-mediated transcription, yet are not growth inhibited by TGF- β 1. We further observed that TGF- β directly contributes to the metastatic behavior of this cell line. Exposure to TGF- β caused 4T1 cells to undergo morphological changes associated with the metastatic phenotype and invade more readily through collagen coated matrices. Furthermore, expression of a dominant negative truncated type II receptor diminished TGF- β signaling and significantly restricted the ability of 4T1 cells to establish distant metastases. Our results suggest that regardless of 4T1 resistance to TGF- β -mediated growth inhibition, TGF- β signaling is required for tumor invasion and metastases formation.

© 2001 Wiley-Liss, Inc.

Key words: TGF- β ; Smad; metastasis; mammary cancer; 4T1

Tumor invasion and metastasis are complex processes in which cancer cells detach from the original tumor mass to establish metastatic foci at distant sites. Metastatic cells characteristically lose growth inhibitory responses, undergo alterations in adhesiveness and demonstrate enhanced production of enzymes that can degrade extracellular matrix components.^{1,2} Since it is the development of metastatic disease that is primarily responsible for cancer mortality, an understanding of the mechanisms that facilitate metastatic tumor progression is of great importance.

One cytokine that may contribute to the metastatic potential of tumor cells is transforming growth factor beta (TGF- β). Originally identified as a positive growth factor for mesenchymal cells, TGF- β has been identified as a potent growth inhibitor of most cell types, including cells of hematopoietic origin and epithelial lineage.³ TGF- β inhibits the growth of normal epithelial cells by inducing an arrest in G1 phase of the cell cycle and, less commonly, by promoting apoptosis.^{3,4} In contrast, most malignant cells are refractory to TGF- β -mediated growth arrest⁵. This loss of sensitivity has been linked to tumor progression, and may be due to loss or mutation of TGF- β receptors or dysregulation of TGF- β signal transduction pathways.^{2,4,6}

It is widely accepted that TGF- β promotes tumorigenicity by stimulating angiogenesis,⁷ inducing extracellular matrix degradation⁸ and inhibiting anti-tumor immune responses.^{9–12} Recent studies have shown that tumorigenicity can also arise via the action of TGF- β on the tumor cells directly.^{12–19} In this report, we have employed a highly metastatic murine mammary cancer cell line (4T1) to examine the role of TGF- β on invasion and metastatic potential *in vivo*. We found that although 4T1 mammary tumor cells are resistant to TGF- β -mediated growth inhibition, TGF- β signaling is critical to tumor invasion and metastases formation. Engagement of ligand initiated downstream signaling pathways

that culminated in Smad2 phosphorylation and transcription of a Smad responsive reporter gene (3TP-lux). Furthermore, disruption of TGF- β signaling by expression of a dominant negative truncated type II receptor significantly curbed tumor metastasis without affecting primary tumor growth.

MATERIAL AND METHODS

Cell lines

The metastatic line 4T1 is a thioguanine-resistant variant of 410.4, a tumor subline isolated from a spontaneous mammary tumor that developed in a BALB/c/c3H mouse. The 4T1 cell line was kindly provided by Dr. Fred Miller of the Michigan Cancer Foundation (Detroit, MI). The TGF- β sensitive Mink lung epithelial cell line (Mv1Lu) was purchased from ATCC (CCL64; Rockville, MD). Both cell lines were maintained *in vitro* by passage in Iscove's Modified Dulbecco's Medium (IMDM) supplemented with 10% fetal bovine serum (FBS).

Mice

Female C.B-17/IcrACCscid mice, 6–8 weeks old, were purchased from a colony maintained at the University of Arizona. The mice were housed in the University of Arizona animal facilities in accordance with the principles of animal care (NIH publication No. 85-23, revised 1985).

Cell cycle analysis

4T1 tumor cells and TGF- β sensitive Mv1Lu cells were arrested at the G1 phase of the cell cycle by seeding them at high density (10^7 cells/T75 flask) for 72 hr. The cells were then detached and plated at a concentration of 5×10^5 cells/100 mm \times 15 mm tissue culture dish in the presence of increasing amounts of TGF- β 1. Forty-eight hours later, the cells were collected, fixed at a concentration of 10^6 cells/ml with cold 70% ethanol for 2 hr and stored at 4°C for up to 1 week. For analysis, 10^6 cells were suspended in 1 ml of Vindelov's PI buffer pH 8.0 (10 mM Trizma base, 10 mM NaCl, 0.1% NP-40, 50 μ g/ml RNase A and 50 μ g/ml propidium iodide). After a 20 min incubation at room temperature, the cells

Grant sponsor: NIH; Grant numbers: CA6929-01A2, CA62212, CA59702; Grant sponsor: Arizona Disease Control Research Commission; Grant number: 9801; Grant sponsor: NIH Institutional Core Grant; Grant number: CA23074; Grant sponsor: NIEHS; Grant number: ES06694; Grant sponsor: Department of Veteran Affairs

*Correspondence to: Dr. Emmanuel T. Akporiaye, University of Arizona, Department of Microbiology and Immunology, 1501 North Campbell Tucson, AZ 85724. Fax: 520 626-2100, E-mail: akporiaye@u.arizona.edu.

Received 9 March 2000; Revised 26 July 2000; Accepted 5 September 2000

were filtered through a 30 μ m nylon mesh and measured for DNA content with a FACStar^{PLUS} flow cytometer (Becton Dickinson Immunocytometry Systems; San Jose, CA) using an Innova 90-5 argon laser (Coherent; Palo Alto, CA) tuned to 488 nm at 100 mW. Fluorescence emission was captured through a 575/26 nm bandpass filter. Data were acquired using Lysys II software and analyzed with CELL-FIT (Becton Dickinson Immunocytometry Systems).

Determination of TGF- β receptor expression

Reverse-transcriptase polymerase chain reaction (rt-pcr). Total RNA was isolated from 4T1 and Mv1Lu cells with TRIzol reagent (GIBCO BRL Life Technologies, Inc., Gaithersburg, MD), treated with DNase I and reverse transcribed to generate cDNA (cDNA cycle kit number K1310-02, Invitrogen; San Diego, CA). PCR amplification was performed in a 50 μ l volume containing 2 μ l of the cDNA reaction, 1.5 mM MgCl₂, 200 μ M dNTPs, 0.1 μ M sense and antisense primers for TGF- β RI or TGF- β RII, and 2.5 units AmpliTaq Gold DNA polymerase (N808-0241, PE Applied Biosystems; Foster City, CA) in 1 \times PCR buffer. Cycling parameters were 95°C for 10 min followed by 94°C for 10 sec, 65°C for 30 sec and 75°C for 45 sec for 50 cycles. TGF- β RI- and TGF- β RII-specific primers were designed using the OLIGO program (National Biosciences; Plymouth, MN) and were as follows: TGF- β RI upper primer (5'-GGGGCGAAGGCATTACAGTG-3', position 76) and TGF- β RI lower primer (5'-ATTGGCACACGGTGGT-GAA-3', position 461); TGF- β RII upper primer (5'-TCCACGT-GCGCCAACAACAT-3', position 915) and TGF- β RII lower primer (5'-GCGCAAGGACAGCCCGAAGT-3', position 1420). These primers amplify 405 bp and 525 bp fragments of TGF- β RI and TGF- β RII, respectively.

¹²⁵I-TGF- β 1 affinity crosslinking studies. Subconfluent 4T1 tumor cells in 12-well plates were labeled with 100 pM ¹²⁵I-TGF- β 1 (173 μ Ci/ μ g, DuPont NEN; Boston, MA) in the presence or absence of 5 nM unlabeled rhTGF- β 1 (Genentech, South San Francisco, CA) for 4 hr at 4°C. After washing, the specifically bound labeled ligand was cross-linked to cell surface receptors with 1 mM bis(sulfosuccinimidyl) suberate (BS³) (Pierce; Rockford, IL) as described previously.²⁰ Equivalent amounts of protein were then resolved by 3–12% gradient SDS-PAGE and labeled receptors visualized by autoradiography.

Cell migration and invasion assays

Cells were seeded at a density of 10⁵ cells per well in triplicate in the upper chamber of 12-well transwells (8 μ m pore, number 29442-120, Corning, Inc., Corning, NY) in the presence or absence of 2 ng/ml rhTGF- β 1 or anti-TGF- β neutralizing antibody (100 μ g/ml) (ATCC HB-9849). Twenty hours later, the transwells were removed and the lower chambers were incubated for an additional 20 hr. Cells that migrated through the transwell into the lower chamber and attached were treated with trypsin and counted. The transwells were untreated (migration assay) or coated with rat tail type I collagen (invasion assay) as previously described.²¹

Confocal microscopy

Cells (10⁴) were grown on 0.17 mm thick coverslips in a CO₂-humidified incubator for 48 hr in the presence or absence of 2 ng/ml rhTGF- β 1. The cells were then fixed with 4% methanol-free formaldehyde for 20 min at room temperature, permeabilized with 100% methanol at -20°C for 6 min, air-dried and stored at -20°C until time of staining. Morphological changes in tumor cells treated with TGF- β were detected by staining the cells with 100 μ l of a 1:40 dilution of bodipy-phalloidin (number B-3416, Molecular Probes, Eugene, OR), a fluorescent molecule that binds to actin filaments. The cells were visualized with a LEICA confocal microscope.

Western blotting

Cells grown overnight in serum free IMDM were untreated or treated with 0.25 ng/ml of rhTGF- β 1 for 45 min, washed twice with ice-cold PBS and scraped into a buffer containing 50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1% NP-40, 0.25% Na-deoxy-

cholate, 1 mM EDTA, 1 mM PMSF, 1 mM NaF, 1 mM Na₃VO₄, 1 μ g/ml aprotinin, 1 μ g/ml leupeptin and 1 μ g/ml pepstatin. The lysate was placed on a rocker at 4°C for 30 min, centrifuged at 14,000g and protein in the supernatant recovered and quantified by BCA Protein Assay (Pierce, Rockford, IL). Proteins (25 μ g) from the cell lysates were resolved by SDS-8% PAGE and electrotransferred to polyvinylidene fluoride (PVDF) membrane. Nonspecific binding sites were saturated by incubation in TBS containing 0.1% Tween-20 and 5% nonfat powdered milk. Membrane bound anti-Smad2 and anti-phosphorylated Smad2 antibodies (S66220; Transduction Laboratories, Lexington, KY and 06-829; Upstate Biotech, Lake Placid NY, respectively) were visualized with horseradish peroxidase-conjugated secondary antibodies (Santa Cruz Biotech) and enhanced chemiluminescence (Amersham Pharmacia Biotech, UK).

Plasmids

A plasmid construct containing cDNA encoding a truncated TGF- β RII lacking the serine/threonine kinase domain (DNRII)²² was kindly provided by Dr. Barbara Ballerman of The Johns Hopkins University. For our studies, the 0.58 kb DNRII fragment was excised by a NheI/Xho digest and cloned into a commercially available plasmid pcDNA3.1zeo(+) (V860-20, Invitrogen). Proper insertion of this fragment into this plasmid was confirmed by restriction analysis and sequencing. The resultant construct (DNRII-pcDNAzeo) contained the truncated receptor under control of human CMV promoter and a SV40-driven zeocin resistance gene to allow for selection of stably transfected cells. The p3TPLux TGF- β -inducible luciferase construct was a gift from Dr. Joan Massagué (Howard Hughes Medical Institute, Memorial Sloan-Kettering Cancer Center, New York, NY) and has been described previously.²³ The β -galactosidase expression vector VR1412 used to normalize luciferase activities was kindly provided by Vical Inc. (San Diego, CA).

Transcriptional response assay

Cells were plated to semiconfluency and 24 hr later were transiently co-transfected with 1 μ g each of p3TPLux and a control vector (VR1412) with Lipofectamine reagent (number 18324-012, GIBCO BRL) according to the manufacturer's instructions. Twenty-four hours after transfection, the cells were plated in triplicate at a concentration of 5 \times 10⁴ cells/well of a white 96-well culture plate (number 6005180, Instrument Company, Inc., Meriden, CT) in the presence or absence of 2 ng/ml rhTGF- β 1 for 18 hr. Extracts were then prepared and assayed for luciferase and β -galactosidase activity using the LucLite Plus (number 6016961, Packard) and Galacto-Light Plus (BL100P, Tropix, Bedford, MA) kits, respectively. Light emission was detected with a Packard Lumicount and luciferase activity was normalized on the basis of β -galactosidase expression.

Northern blot analysis

Messenger RNA was isolated from subconfluent cells and subjected to Northern blot analysis using a NorthernMax-Gly kit (number 1944; Ambion, Inc., Austin, TX). Briefly, the mRNA was separated on a 1% agarose gel, transferred to nylon membrane, UV crosslinked and hybridized with biotin-labeled rat TGF- β receptor II and mouse GAPDH-specific RNA probes overnight at 80°C and 65°C, respectively. Hybridized bands were visualized by the addition of streptavidin/alkaline phosphatase using a BrightStar Bio-detect kit (number 1925; Ambion, Inc.). To develop an antisense TGF- β receptor II (T β RII) template for transcription of a T β RII-specific probe, the DNRII fragment obtained from a NheI/XhoI digest was cloned into pcDNA3.1(+) that had been digested with XhoI and XbaI. To generate the sense T β RII RNA probe, this template was linearized by ApaI digestion, treated with proteinase K and purified on a 4% polyacrylamide gel. The linearized plasmid was transcribed with T7 polymerase in the presence of a nucleotide triphosphate RNA-labeling mix (number 1685 597; Roche Molecular Biochemicals, Indianapolis, IN) containing biotin-16-UTP

according to the manufacturer's instructions. Biotin-labeled RNA probe specific for GAPDH and RNA millennium markers were similarly generated from linearized templates (number 7431 and number 7785, respectively; Ambion, Inc.).

Spontaneous metastasis assay

Mice challenged orthotopically in the mammary gland with 10^5 cells were sacrificed 21 days post-injection. Lungs were removed, stained with India ink and the tumors bleached with Fekete's solution as previously described.²⁴ Primary tumors were measured in 2 dimensions every 3–4 days post inoculation. Tumor volume was calculated using the formula $v = (l \times w^2)/2$, where v = volume (mm^3), l = long diameter and w = short diameter²⁴.

RESULTS

TGF- β does not inhibit cell cycle progression of 4T1 cells

We assessed whether exposure of 4T1 tumor cells to TGF- β resulted in growth inhibition. For this purpose, cells were arrested at G1 by crowding, released to transit through the cell cycle in the presence or absence of increasing amounts of TGF- β and then analyzed by flow cytometry. As shown in Figure 1, 4T1 cells were not growth arrested in G1 but instead progressed through the cell cycle in the presence of high concentrations (20 ng/ml) of TGF- β 1. In contrast, indicator Mink lung epithelial cells (Mv1Lu) that are growth inhibited by TGF- β were arrested in G1 following exposure to 2 ng/ml TGF- β 1 (Fig. 1). The resistance of 4T1 cells to TGF- β growth control was additionally supported by cell counting; no differences in cell number and viability were observed between untreated ($2.3 \times 10^6 \pm 3.6 \times 10^5$) and TGF- β -treated cells ($2.4 \times 10^6 \pm 1.5 \times 10^4$) after 48 hr of *in vitro* culture.

Expression of functional TGF- β receptors in mammary tumor cells

Since 4T1 cells were resistant to TGF- β -mediated cell cycle control, we examined them for the presence of functional TGF- β receptors. Analysis of RNA by RT-PCR demonstrated that 4T1 cells transcribe mRNA for TGF- β receptors I and II (Fig. 2a). The PCR product obtained for TGF- β RI was sequenced and determined to be ALK-5 (data not shown). 4T1 cells also express TGF- β RI and TGF- β RII as surface proteins capable of binding TGF- β ligand. When labeled with ^{125}I -TGF- β 1 in a receptor cross-linking experiment, TGF- β receptors I, II and III were detected in 4T1 cells (Fig. 2b). Binding of ^{125}I -TGF- β 1 was com-

petitively inhibited by the addition of an excess of unlabeled TGF- β 1 (Fig. 2b).

TGF- β signaling in 4T1 tumor cells

Next we evaluated if TGF- β signaling occurred in 4T1 tumor cells following TGF- β treatment. The involvement of Smads in TGF- β signaling was determined by Western blot to detect the presence of phosphorylated Smad-2 and by gene expression analysis of cells transiently transfected with the Smad responsive 3TP-Lux reporter construct. Smad2 phosphorylation and gene transcription in unstimulated 4T1 cells were significantly increased upon addition of exogenous TGF- β 1 (Fig. 3).

Morphological changes induced by TGF- β treatment

The 4T1 mammary tumor cell line typically adheres to plastic or glass substratum and grows in rounded clumps (Fig. 4a). When incubated with 2 ng/ml rhTGF- β 1, the cells became less clumped, assumed a fibroblast-like appearance and demonstrated reorganization of the cytoskeleton characterized by prominent F-actin filaments and stress fibers (Fig. 4b). In order to determine if this morphologic change involved an epithelial to mesenchymal transition (EMT), cells were stained with vimentin and keratin antibodies as previously described.²⁵ 4T1 cells constitutively co-expressed vimentin and keratin intermediate filaments and TGF- β treatment did not alter their expression (data not shown). This interconverted phenotype has been previously reported and is characteristic of highly malignant breast cancer cells.^{25,26–28}

TGF- β stimulates *in vitro* migration and invasion of 4T1 cells

Since SMAD-mediated-TGF- β signaling was intact in 4T1 cells and TGF- β treatment induces morphologic changes, we speculated that TGF- β might contribute to tumor progression by stimulating cell migration and invasiveness. To explore this possibility, the effect of TGF- β on *in vitro* migration and invasion of 4T1 cells was evaluated. The data (Fig. 5) demonstrated basal levels of migration but negligible invasive capacity in 4T1 cells in the absence of TGF- β . Addition of TGF- β resulted in a 4-fold increase in *in vitro* migration and a 17-fold increase in invasiveness that was completely abrogated in the presence of TGF- β neutralizing antibody (Fig. 5).

Disruption of TGF- β signaling inhibits metastasis formation

To directly evaluate the role of TGF- β signaling in tumorigenicity and the development of metastasis, 4T1 cells were trans-




FIGURE 1 – Effect of TGF- β on the cell cycle of 4T1 cells. Mv1Lu and 4T1 cells arrested in G1 by crowding were released from growth arrest by detaching and plating at a lower cell density in the presence of different concentrations of rhTGF- β 1. After 48 hr, the cells were collected and evaluated for DNA content by flow cytometric analysis as described in Material and Methods.




FIGURE 2 – Expression of TGF- β receptors by 4T1 tumor cells. (a) RT-PCR was performed on total RNA isolated from 4T1 (lane 2) and Mv1Lu cells (lane 4) using primers specific for a 405 bp fragment of TGF- β R1, a 525 bp fragment of TGF- β R2 or a 215 bp fragment of histone. Lanes 1, 3 and 5 represent PCR reactions containing no template, 4T1 RNA or Mv1Lu RNA (no RT) respectively. (b) Subconfluent 4T1 tumor cells were labeled with 100 pM 125 I-TGF- β 1 in the presence (+) or absence (–) of 5 nM competing unlabeled TGF- β 1. Bound labeled ligand was cross-linked to cell surface receptors, the proteins resolved by SDS-PAGE and labeled receptors visualized by autoradiography.




FIGURE 3 – TGF- β induction of Smad2 phosphorylation and transcriptional responses in 4T1 cells. A. Cells were incubated in the presence (+) or absence (–) of 2 ng/ml rhTGF- β for 45 min. Cell lysates were separated by SDS-PAGE, transferred to membrane and incubated with pretested dilutions of antibodies specific for Smad2 or the phosphorylated form of Smad2 (55 kDa). (b) 4T1 cells cotransfected with p3TP-lux reporter and VR1412 β -galactosidase control plasmids were treated with 2 ng/ml rhTGF- β 1 and/or anti-TGF- β neutralizing antibody 18 hr before determination of luciferase activity. Bars represent the mean relative light units and standard deviation of triplicate samples. Luciferase activities were normalized based on β -galactosidase expression.

fects with a truncated dominant-negative TGF- β type II receptor (DNR2). This receptor lacks the cytoplasmic serine/threonine kinase domain and is therefore unable to transduce TGF- β -initiated

signals²². Expression of DNR2 was demonstrated by Northern blot (Fig. 6) and 4 of the stably transfected clones were selected for further analysis. Expression of DNR2 blunted tran-




FIGURE 4 – Effect of TGF- β on tumor cell morphology. Subconfluent 4T1 cells were grown on glass coverslips for 48h in culture medium in the absence (a) or presence (b) of rhTGF- β 1. Cells were visualized with a Lieca confocal microscope after staining actin fibers with bodipy-phalloidin. Magnification = 400 \times .




FIGURE 5 – TGF- β 1 enhances migration and invasion of 4T1 cells. Tumor cells were placed in the upper chamber of 12-well transwells and incubated in the presence of anti-TGF- β antibody, rat IgG (isotype) or rhTGF- β 1. Cells that moved through untreated transwells (migration assay) or collagen coated transwells (invasion assay) into the lower chamber were counted. Bars represent the mean cell number \pm SE of duplicate wells. The data shown are representative of at least 2 experiments.

scriptional activation by TGF- β in the majority of clones tested (Fig. 7). Additionally, the expression of DNRII resulted in a decrease of TGF- β -induced phosphorylation of Smad2 (Fig. 7b). Furthermore, when injected orthotopically into the mammary gland of immunodeficient SCID mice, 3 of the 4 clones expressing the DNRII were severely inhibited in their ability to form metastatic lung nodules (Table I). Approximately 114 ± 35 (n=4) metastatic nodules were found in the lungs of mice that were orthotopically injected with mock-transfected 4T1 cells (4T1Zeo) compared with 15.5 ± 6.5 (n=15) in mice injected with DNRII-expressing tumor cells. In the majority of mice injected with gene-modified tumor cells, the sizes of the primary tumors were not significantly different from those in control animals injected with tumor cells transfected with vector alone (Table I). Histolog-




FIGURE 6 – Expression of dominant negative truncated TGF- β receptor II in transfected 4T1 cells. mRNAs isolated from mock-transfected control cells (Zeo) and 4T1 cells stably transfected with DNRII-pcDNAzeo (clones 5, 25, 32 and 34) were subjected to Northern blot hybridization with TGF- β receptor II and GAPDH-specific RNA probes.

ically, no discernible differences in the morphology or local invasiveness of the primary tumors were observed among the test and control groups. All primary tumors were well demarcated and extended from the epidermal/dermal junction through the dermis with infiltration into the abdominal musculature. Each tumor was bordered by mild neutrophilic inflammatory infiltrates and contained extensive necrotic foci. Carcinoma cells were not observed in blood vessels adjacent to the primary tumors. In contrast to the effects on *in vivo* metastasis, the expression of the truncated TGF- β type II receptor did not affect *in vitro* migration and invasion (data not shown).

DISCUSSION

In our study we investigated the role of TGF- β in the metastatic phenotype of a highly metastatic murine 4T1 mammary tumor. We found that TGF- β contributes to the ability of the tumor cells to




FIGURE 7 – Gene transcription in dominant negative TGF- β receptor II expressing clones. (a) 4T1 cells stably transfected with a mock vector (Zeo) or the dominant negative truncated TGF- β receptor II (clones 5, 25, 32 and 34) were transiently co-transfected with p3TP-lux reporter and VR1412 β -galactosidase plasmid DNA. Cells were then treated with rhTGF- β 1 for an additional 18 hr before determination of luciferase activity. Bars represent fold increase and standard error of triplicate samples. Luciferase activities were normalized based on β -galactosidase expression. The data shown are representative of 3 experiments. (b) 4T1 cells stably transfected with a mock vector (Zeo) or the dominant negative truncated TGF- β receptor II (clones 5, 25, 32 and 34) were treated with rhTGF- β 1 and immunoblotting for Smad2 and the phosphorylated form of Smad2 was performed.

TABLE I – EXPRESSION OF TRUNCATED TGF- β RECEPTOR II SUPPRESSES METASTASES FORMATION¹

Cell type	Tumor volume	Number of lung metastases
4T1	783.4 \pm 34.2	251 \pm 36
Zeo	810.7 \pm 102.8	114 \pm 35
clone 5	899.4 \pm 110.6	10 \pm 3
clone 25	606.7 \pm 53.2	1 \pm 1
clone 32	952.1 \pm 192.4	61 \pm 13
clone 34	407.2 \pm 23.9	3 \pm 1

¹SCID mice were injected orthotopically with parental, mock transfected or gene-modified 4T1 cells. Three weeks post-injection, the mice were sacrificed and their lungs were collected and examined for the presence of surface metastases. Values represent the mean \pm standard error (n = 4).

invade and metastasize *in vivo*. The results also exhibit that 4T1 cells are resistant to TGF- β -mediated growth inhibition yet respond to TGF- β as demonstrated by the presence of membrane receptors that bind ligand, initiate downstream phosphorylation of Smad2 and induce Smad-dependent transcriptional activation. Thus, unlike numerous cancer cells whose resistance to TGF- β growth inhibition is due to inactivation of TGF- β receptors^{29–31} or Smad family signal transducers,³² 4T1 cells are capable of TGF- β -receptor-mediated signal transduction. These findings are in concordance with those of others that have identified functional receptors in TGF- β growth resistant cancers such as glioma³³, melanoma³⁴ and cervical carcinoma.³⁵ The intactness of Smad-mediated gene transcription in the 4T1 mammary tumor cell line is similar to that reported in human melanoma cells in which Smad-mediated gene transcription was shown to be independent of cell proliferation.³⁴

The TGF- β -induced morphological changes that we observed in 4T1 mammary tumor cells are consistent with those reported by others in transformed murine mammary epithelial cells,¹⁵ murine colon cancer cells¹⁴ and skin carcinoma cells¹⁶ and have been

correlated with metastatic potential.^{14–16} Recent work by Piek et al.¹⁸ suggests that Smad proteins may play a role in the invasive and metastatic phenotype. Using the NMuMG murine mammary epithelial cell line they demonstrated the requirement for Smad2 and Smad4 proteins in inducing transdifferentiation in cells constitutively expressing low levels of type I receptor (ALK-5). The transition from the epithelial to the mesenchymal phenotype was characterized by stress fiber formation and downregulation/relocalization of E-cadherin, changes frequently associated with the invasive and metastatic phenotype^{14,15}. Similar Smad-dependent pathways present in 4T1 tumor cells may contribute to their invasive and metastatic activities. That 4T1 cells secrete TGF- β and exhibit basal levels of Smad2 phosphorylation suggests a cell autonomous mechanism of TGF- β signaling which may contribute to the metastatic phenotype as has been proposed in murine models of renal, lung, mammary and colon cancers.^{14,17,19} Experiments are in progress to resolve this question.

The importance of TGF- β in the invasion and metastasis of 4T1 tumor cells was demonstrated in studies in which expression of a truncated dominant negative receptor significantly suppressed formation of lung metastases *in vivo*. Whereas metastasis formation was severely curbed, tumorigenicity at the primary tumor injection site was not affected. This observation is in agreement with that of Yin et al.¹⁹ using the human breast cancer cell line MDA-MB-231 in which they exhibited that dominant-negative blockade of the type II TGF- β receptor had no effect on the growth of local tumor yet significantly diminished metastasis to the bone.¹⁹ In our study, histological analysis did not reveal any discernible differences in the morphology or local invasiveness of the primary tumors between mock-transfected (4T1-Zeo) and DNRII-expressing tumors. In addition, no histologic evidence of hematogenous metastases was observed in blood vessels adjacent to the primary tumors. These findings are not unexpected since hematogenous invasion of neoplastic cells is a relatively infrequent occurrence that is rarely documented histologically. A more reliable indicator of hematogenous invasion of carcinoma cells is the presence of metastases at

distant sites such as the lung. The decreased number of lung metastases in the DNR11 group is therefore considered a significant finding in our study and reflective of decreased hematogenous invasion of the carcinoma cells that express the mutant receptor. Taken together, our findings suggest that TGF- β may preferentially exert its effect on cellular genes that control invasion and spread of tumor cells.

Our observation that the DNR11 affected *in vivo* metastasis but *not in vitro* migration and invasion suggests that *in vitro* assays are limited in their ability to predict metastatic capacity and that characteristics other than increased motility are required for spread to occur *in vivo*. TGF- β may promote metastasis in part through its ability to regulate the expression or activity of extracellular matrix-degrading proteases like MMP-9 whose role in tumor metastasis is well established.^{36,37} Our finding that TGF- β upregulates the se-

cretion of MMP-9 by 4T1 cells (unpublished observation) is in agreement with this mechanism. Understanding the TGF- β -induced downstream effectors mediating the metastatic phenotype will likely lead to the identification of molecular targets that once perturbed could override this phenotype.

ACKNOWLEDGEMENTS

The authors thank Barbara Carolus for expert technical assistance. This work was supported by NIH Grants CA6929-01A2 (E.T.A.), CA59702 (M.J.C.H.) the Arizona Disease Control Research Commission (contract #9801)(E.T.A.), NIH Institutional Core Grants #CA23074, NIEHS Grant # ES06694, NIH grant CA62212 (C.L.A.), and a Clinical Investigator Award from the Department of Veteran Affairs (C.L.A.).

REFERENCES

- Woodhouse EC, Chuaqui RF, Liotta LA. General mechanisms of metastasis. *Cancer* 1997;80:1529-37.
- Reiss M, Barcellos-Hoff MH. Transforming growth factor- β in breast cancer: a working hypothesis. *Breast Cancer Res Treat* 1997;45:81-95.
- Taipale J, Saharinen J, Keski-Oja J. Extracellular matrix-associated transforming growth factor- β : role in cancer cell growth and invasion. *Adv Cancer Res* 1998;75:87-134.
- Alexandrow MG, Moses HL. Transforming growth factor β and cell cycle regulation. *Cancer Res* 1995;55:1452-57.
- Fynan TM, Reiss M. Resistance to inhibition of cell growth by transforming growth factor-beta and its role in oncogenesis. *Crit Rev Oncol* 1993;4:493-540.
- Yingling JM, Wang X-F, Bassing CH. Signaling by the transforming growth factor- β receptors. *Biochim Biophys Acta* 1995;1242:115-36.
- Roberts AB, et al. Transforming growth factor type-beta: Rapid induction of fibrosis and angiogenesis in vivo and stimulation of collagen formation in vitro. *Proc Natl Acad Sci USA* 1986;83:4167-71.
- Albo D, Berger DH, Wang TN, Hu X, Rothman V, Tuszynski GP. Thrombospondin-1 and transforming growth factor-beta promote breast tumor cell invasion through up-regulation of the plasminogen/plasmin system. *Surgery* 1997;122:493-500.
- Torre-Amione G, et al. A highly immunogenic tumor transfected with a murine transforming growth factor type β 1 cDNA escapes immune surveillance. *Proc Natl Acad Sci USA* 1990;87:1486-90.
- Arteaga CL, Hurd SD, Winnier AR, Johnson MD, Fendly BM, Forbes JT. Anti-transforming growth factor (TGF)- β antibodies inhibit breast cancer cell tumorigenicity and increase mouse spleen natural killer cell activity. Implications for a possible role of tumor cell/host TGF- β interactions in human breast cancer progression. *Clin Invest* 1993;92:2569-76.
- Park JA, Wang E, Kurt RA, Schluter SF, Hersch EM, Akporiaye ET. Expression of an antisense transforming growth factor- β 1 transgene reduces tumorigenicity of EMT6 mammary tumor cells. *Cancer Gene Ther* 1997;4:42-50.
- Wojtowicz-Praga S. Reversal of tumor-induced immunosuppression: a new approach to cancer therapy. *J Immunother* 1997;20:165-177.
- Miettinen PJ, Ebner R, Lopez AR, Derynck R. TGF- β induced transdifferentiation of mammary epithelial cells to mesenchymal cells: involvement of type I receptors. *J Cell Biol* 1994;127:2021-36.
- Oft M, Heider K-H, Beug H. TGF β signaling is necessary for carcinoma cell invasiveness and metastasis. *Curr Biol* 1998;8:1243-52.
- Oft M, Peli J, Rudaz C, Schwarz H, Beug H, Reichmann E. TGF- β 1 and Ha-Ras collaborate in modulating the phenotypic plasticity and invasiveness of epithelial tumor cells. *Genes Dev* 1996;10:2462-77.
- Portella G, et al. Transforming growth factor β is essential for spindle cell conversion of mouse skin carcinoma in vivo: implications for tumor invasion. *Cell Growth Diff* 1998;9:393-404.
- Hojo M, et al. Cyclosporine induces cancer progression by a cell-autonomous mechanism. *Nature* 1999;397:530-4.
- Piek E, Moustakas A, Kurisaki A, Heldin C-H, Ten Dijke P. TGF- β type I receptor/ALK-5 and Smad proteins mediate epithelial to mesenchymal transdifferentiation in NMuMG breast epithelial cells. *J Cell Sci* 1999;112:4557-68.
- Yin J et al. TGF- β signaling blockade inhibits PTHrP secretion by breast cancer cells and bone metastasis development. *J Clin Invest* 1999;103:197-206.
- Dumont N, O'Connor-McCourt MD, Philip A. Transforming growth factor-beta receptors on human endometrial cells; identification of the type I, II, and III receptors and glycosyl-phosphatidylinositol anchored TGF-beta binding proteins. *Mol Cell Endocrinol* 1995;111:57-66.
- Keely PJ, Westsick JK, Whitehead IP, Der CJ, Parise LV. Cdc42 and Rac1 induce integrin-mediated cell motility and invasiveness through PI(3)K. *Nature* 1997;390:632-6.
- Choi ME, Ballerman BJ. Inhibition of capillary morphogenesis and associated apoptosis by dominant negative mutant transforming growth factor- β receptors. *J Biol Chem* 1995;270:21144-50.
- Wrana JL et al. TGF- β signals through a heteromeric protein kinase receptor complex. *Cell* 1992;71:1003-14.
- Wexler H. Accurate identification of experimental pulmonary metastases. *J Natl Cancer Inst* 1966;36:641-5.
- Hendrix MJC, Seftor EA, Seftor REB, Trevor KT. Experimental co-expression of vimentin and keratin intermediate filaments in human breast cancer cells results in phenotypic interconversion and increased invasive behavior. *Am J Pathol* 1997;150:483-95.
- Thompson EW, et al. Association of increased basement membrane invasiveness with absence of estrogen receptor and expression of vimentin in human breast cancer cell lines. *J Cell Physiol* 1992;150:534-44.
- Sommers CL, et al. Loss of epithelial markers and acquisition of vimentin expression in adriamycin- and vinblastine-resistant human breast cancer cell lines. *Cancer Res* 1992;52:5190-7.
- Lichtner RB, Julian JA, North SM, Glasser SR, Nicolson GL. Coexpression of cytokeratins characteristic for myoepithelial and luminal cell lineages in rat 13762NF mammary adenocarcinoma tumors and their spontaneous metastases. *Cancer Res* 1991;51:5943-50.
- Laiho M, Weis MB, Massagué J. Concomitant loss of transforming growth factor (TGF)- β receptor types I and II in TGF- β -resistant cell mutants implicates both receptor types in signal transduction. *J Biol Chem* 1990;265:18518-24.
- Park K, et al. Genetic changes in the transforming growth factor β (TGF- β) type II receptor gene in human gastric cancer cells: Correlation with sensitivity to growth inhibition by TGF- β . *Proc Natl Acad Sci USA* 1994;91:8772-6.
- Myeroff LL, et al. A transforming growth factor β receptor type II gene mutation common in colon and gastric but rare in endometrial cancers with microsatellite instability. *Cancer Res* 1995;55:5545-7.
- Zang Y, Musci T, Derynck R. The tumor suppressor Smad4/DPC4 as a central mediator of Smad function. *Curr Biol* 1997;7:270-6.
- Isoe S, et al. Resistance to growth inhibition by transforming growth factor- β in malignant glioma cells with functional receptors. *J Neurosurg* 1998;88:529-34.
- Rodeck U, Nishiyama T, Mauviel A. Independent regulation of growth and Smad-mediated transcription by transforming growth factor β in human melanoma cells. *Cancer Res* 1999;59:547-50.
- Kang SH, et al. Genetic integrity of transforming growth factor beta (TGF-beta) receptors in cervical carcinoma cell lines: loss of growth sensitivity but conserved transcriptional response to TGF-beta. *Int J Cancer* 1998;77:620-5.
- Bernhard EJ, Gruber SB, Muschel RJ. Direct evidence linking expression of matrix metalloproteinase 9 (92-kDa gelatinase/collagenase) to the metastatic phenotype in transformed rat embryo cells. *Proc Natl Acad Sci USA* 1994;91:4293-7.
- Ray JM, Stetler-Stevenson WG. The role of matrix metalloproteinases and their inhibitors in tumour invasion, metastasis and angiogenesis. *Eur Respir J* 1994;7:2062-72.

Autocrine Transforming Growth Factor- β Signaling in Mammary Tumor Cell Invasiveness. Nancy Dumont and Carlos L. Arteaga, Departments of Cell Biology and Medicine, Vanderbilt University School of Medicine and Vanderbilt-Ingram Cancer Center.

Transforming growth factor- β (TGF β) is a pleiotropic growth factor, which plays a critical role in modulating cell growth, differentiation and plasticity. There is increasing evidence that after cells lose their sensitivity to TGF β -mediated growth inhibition, autocrine TGF β signaling may potentially promote tumor progression. Studies have shown that introduction of a dominant negative TGF β type II receptor (T β RII) in a variety of tumor cells can delay tumor formation and prevent the conversion of cells from an epithelial to a more invasive and metastatic mesenchymal phenotype. The purpose of this study is to identify the biochemical pathways perturbed by a dominant negative T β RII that are causally associated with the metastatic phenotype of tumors. In order to do that, the MDA-MB-231 human breast cancer cell line was stably transfected with either a kinase-dead T β RII-K277R (dnT β RII) construct or the vector control. Affinity labeling of cell surface receptors with 125 I-TGF β 1 revealed an increase in the labeling of T β RII on the surface of cells transfected with dnT β RII compared to the vector control or parental cells, suggesting that the dnT β RII was indeed expressed. This was confirmed by immunoprecipitating the affinity-labeled exogenous receptor via its HA tag. The function of the transgene was evaluated by i) examining phosphorylation and nuclear translocation of Smad2, a TGF β signal transducer, and ii) measuring transcription utilizing the TGF β responsive promoters, p3TP-lux and p(CAGA) $_{12}$ -lux. These assays revealed that TGF β -mediated Smad2 phosphorylation, Smad2 nuclear translocation, and transcriptional responses were reduced in cells stably transfected with dnT β RII compared to vector control. In addition, wound closure assays indicated that the basal migratory potential of dnT β RII expressing cells was impaired. Biochemical experiments are currently underway to identify the signal transducers perturbed by the dnT β RII expression which impair motility, and which may, in turn, be relevant to TGF β -mediated invasion and metastases.

ABSTRACT GORDON CONFERENCE

Expression of a Kinase Inactive TGF β Type II Receptor Impairs Motility in Breast Cancer Cells

Nancy Dumont, Andrei Bakin and Carlos L. Arteaga, Departments of Cancer Biology and Medicine, Vanderbilt University School of Medicine and Vanderbilt-Ingram Cancer Center.

Transforming growth factor- β (TGF β) is a pleiotropic growth factor, which plays a critical role in modulating cell growth, differentiation and plasticity. There is increasing evidence that after cells lose their sensitivity to TGF β -mediated growth inhibition, autocrine TGF β signaling may potentially promote tumor progression. Studies have shown that introduction of a dominant negative TGF β type II receptor (T β RII) in a variety of tumor cells can delay tumor formation and prevent the conversion of cells from an epithelial to a more invasive and metastatic mesenchymal phenotype. The purpose of this study is to identify the biochemical pathways perturbed by a dominant negative T β RII that are causally associated with the metastatic phenotype of tumors. To accomplish this aim, the MDA-MB-231 metastatic human breast cancer cell line was stably transfected with either a kinase-dead T β RII-K277R (dnT β RII) construct or the vector control. Affinity labeling of cell surface receptors with 125 I-TGF β 1 revealed an increase in the labeling of T β RII on the surface of cells transfected with dnT β RII compared to the vector control or parental cells, suggesting that the dnT β RII was expressed. This was confirmed by immunoprecipitating the affinity-labeled exogenous receptor via its HA tag. The function of the transgene was evaluated by i) examining phosphorylation and nuclear translocation of Smad2, a TGF β signal transducer, and ii) measuring transcription utilizing the TGF β responsive promoters, p3TP-lux and p(CAGA) $_{12}$ -lux. These assays revealed that TGF β -mediated Smad2 phosphorylation, Smad2 nuclear translocation, and transcriptional responses were reduced in cells stably transfected with dnT β RII compared to vector control. In addition, wound closure assays indicated that the basal migratory potential of dnT β RII expressing cells was impaired, independent of changes in proliferation. This impairment in motility appears to be TGF β specific in that it can be restored by expression of a constitutively active type I TGF β receptor, but not by constitutively active activin or BMP type I receptors. Biochemical experiments are currently underway to identify the signal transducers perturbed by dnT β RII expression which impair motility, and which may, in turn, be relevant to TGF β -mediated invasion and metastases.